

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00852 A2

(51) International Patent Classification⁷: C12N 9/00

Hjortespringparken 40, DK-2730 Herlev (DK). VIND, Jesper [DK/DK]; Bagsværdvej 115, DK-2800 Lyngby (DK).

(21) International Application Number: PCT/DK01/00448

(22) International Filing Date: 26 June 2001 (26.06.2001)

(74) Common Representative: NOVOZYMES A/S; Patents, Krogshøjvej 36, DK-2880 Bagsværd (DK).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

PA 2000 00989	26 June 2000 (26.06.2000)	DK
PA 2000 00990	26 June 2000 (26.06.2000)	DK
PA 2000 00991	26 June 2000 (26.06.2000)	DK
PA 2000 01172	4 August 2000 (04.08.2000)	DK
09/703,416	31 October 2000 (31.10.2000)	US
09/703,414	31 October 2000 (31.10.2000)	US
PA 2001 00843	25 May 2001 (25.05.2001)	DK

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): NOVOZYMES A/S [DK/DK]; Krogshøjvej 36, DK-2880 Bagsværd (DK). NOVOZYMES BIOTECH, INC. [US/US]; 1445 Drew Avenue, Davis, CA 95616 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): TSUTSUMI, Noriko [JP/JP]; 3-2-16, Higashisugano, Ishikawa, Chiba 272 (JP). SASAKI, Yukiko [JP/JP]; 2-26-1 Kitakata, Ishikawa, Chiba 273-0035 (JP). REY, Michael, W. [US/US]; 605 Robin Place, Davis, CA 95616 (US). ZARETSKY, Elizabeth [US/US]; 1114 Colina Ct., Davis, CA 95616 (US). SPENDLER, Tina [DK/DK];

Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: LIPOLYTIC ENZYME

(57) Abstract: A group of genes encoding lipolytic enzymes with high homology have been isolated from strains of *Fusarium* and *Acremonium*.

WO 02/00852 A2

LIPOLYTIC ENZYME

FIELD OF THE INVENTION

The present invention relates to a polynucleotide encoding a lipolytic enzyme and to a lipolytic enzyme. The invention also relates to methods of
5 producing and using the lipolytic enzyme.

BACKGROUND OF THE INVENTION

Lipolytic enzymes (such as lipases and phospholipases) are capable of hydrolyzing carboxylic ester bonds in a substrate to release carboxylic acids. They are known to be useful, e.g., in baking and detergents.

10 Various species of the related genera *Acremonium* and *Fusarium* are known to produce lipolytic enzymes. Thus, Roberts et al., *Mycologia*, 79, 265-273, 1987. Satyanarayana & Johri, *Current Science*, 50, 680-682, 1981 state that some isolates of *Acremonium* produce lipase.

WO 98/26057 discloses a polypeptide having lipase and phospholipase
15 activity (GenBank Acc. No. A85215) obtained from *Fusarium oxysporum*. A lipase from *Fusarium heterosporum* and its sequence are known. *Journal of Fermentation and Bioengineering*, 75 (5), p 349-352, 1993. *Journal of Biochemistry (Tokyo)*, 116 (3), p 536-540, 1994. A lipolytic enzyme from *Fusarium culmorum* CBS 513.94 and its N-terminal sequence are disclosed in US 5830736. A lipase/phospholipase from
20 *Fusarium oxysporum* and its sequence are disclosed in WO 98/26057. US 5990069 discloses a lipase from a strain of *Fusarium solani* var. *minus*.

SUMMARY OF THE INVENTION

The inventors have isolated a group of genes encoding lipolytic enzymes with high homology from *Fusarium* and *Acremonium*.

25 Accordingly, one aspect of the invention provides a polynucleotide encoding a lipolytic enzyme which comprises:

- a) the DNA sequence encoding a mature lipolytic enzyme cloned into a plasmid present in *Escherichia coli* DSM 13539,
- b) the mature polypeptide-encoding DNA sequence shown in SEQ ID NO: 1,
30 3, 5, 7 or 9,
- c) an analogue of the DNA sequence defined in a) or b) which
 - i) has at least 80 % identity with said DNA sequence, or
 - ii) hybridizes at high stringency with said DNA sequence, its complementary strand or a subsequence thereof.

Another aspect of the invention also provides a lipolytic enzyme which may be a polypeptide having an amino acid sequence as the mature peptide shown in SEQ ID NO: 2, 4, 6, 8 or 10. It may be C-terminally processed to consist of amino acids 1-282 of SEQ ID NO: 2, 1-282 of SEQ ID NO: 4, 1-275 of SEQ ID NO: 6, 1-282 of SEQ ID NO: 8 or 1-276 of SEQ ID NO: 10.

Further, the lipolytic enzyme of the invention may be a polypeptide encoded by the lipolytic enzyme encoding part of the DNA sequence cloned into a plasmid present in *Escherichia coli* deposit number DSM 13539.

The lipolytic enzyme may also be an analogue of the polypeptide defined above which:

- i) has at least 85 % identity with said polypeptide,
- ii) is immunologically reactive with an antibody raised against said polypeptide in purified form,
- iii) is an allelic variant of said polypeptide,

Finally, the lipolytic enzyme of the invention may be a polypeptide which is encoded by a nucleic acid sequence which hybridizes under high stringency conditions with a complementary strand of the mature polypeptide-encoding nucleic acid sequence of SEQ ID NO: 1, 3, 5, 7 or 9 or a subsequence thereof having at least 100 nucleotides.

A further aspect of the invention provides a nucleic acid sequence comprising a nucleic acid sequence which encodes the lipolytic enzyme described above.

Other aspects of the invention provide a recombinant expression vector comprising the DNA sequence, and a cell transformed with the DNA sequence or the recombinant expression vector.

The DNA sequences of the invention have the following identities to known DNA sequences:

SEQ ID NO:	Source organism	Closest known sequence	DNA identity
1	<i>F. venenatum</i>	<i>F. oxysporum</i>	79 %
3	<i>F. sulphureum</i>	<i>F. oxysporum</i>	77 %
5	<i>A. berkeleyanum</i>	<i>F. heterosporum</i>	68 %
7	<i>F. culmorum</i>	<i>F. oxysporum</i>	79 %
9	<i>F. solani</i>	<i>F. oxysporum</i>	68 %

DETAILED DESCRIPTION OF THE INVENTION**Genomic DNA source**

A lipolytic enzyme of the invention may be derived from a strain of *Fusarium* or *Acremonium*, particularly *F. venenatum*, *F. sulphureum*, *A. berkeleyanum*, *F. culmorum* or *F. solani*, using probes designed on the basis of the DNA sequences in this specification, or from a Plasmid contained in an *E. coli* strain identified below.

More particularly, the *Fusarium venenatum* cell may be *Fusarium venenatum* A3/5, which was originally deposited as *Fusarium graminearum* ATCC 20334 and recently reclassified as *Fusarium venenatum* by Yoder and Christianson, 1998, *Fungal Genetics and Biology* 23: 62-80 and O'Donnell *et al.*, 1998, *Fungal Genetics and Biology* 23: 57-67. Alternatively, the *Fusarium venenatum* cell may be a morphological mutant of *Fusarium venenatum* A3/5 or *Fusarium venenatum* ATCC 20334, as disclosed in WO 97/26330. ATCC 20334 is available on commercial terms from American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 ("ATCC").

Other strains are *A. berkeleyanum* CBS 301.38, *F. culmorum* CBS 513.94 and *F. solani* MUCL 38667. The two CBS strains are available on commercial terms from Centraalbureau voor Schimmelcultures, Uppsalalaan 8, 3584 CT Utrecht, the Netherlands (P.O.Box 85167, 3508 AD Utrecht, the Netherlands). MUCL 38667 is available on commercial terms from Mycothèque de l'Université Catholique de Louvain, Place Croix du Sud 3, B-1348 Louvain-la-Neuve, Belgium by referring to US 5990069.

Strains of *Escherichia coli* each containing a gene encoding a lipolytic enzyme from a source organism were deposited by the inventors under the terms of the Budapest Treaty as follows:

Source organism	<i>E. coli</i> deposit date	<i>E. coli</i> deposit number
<i>F. venenatum</i> WTY700 3.8d	22 August 2000 (Resubmitted on 30 March 2001)	NRRL B-30333
<i>F. sulphureum</i>	15 June 2000	DSM 13539
<i>A. berkeleyanum</i> CBS 301.38	15 June 2000	DSM 13538
<i>F. culmorum</i> CBS 513.94	15 June 2000	DSM 13537
<i>Fusarium solani</i> MUCL 38667	21 June 2001	DSM 14361

NRRL indicates a deposit made at Agricultural Research Service Patent Culture Collection, Northern Regional Research Center, 1815 University Street, Peoria, Illinois, 61604. NRRL B-30333 was resubmitted on 30 March 2001. DSM indicates a deposit made at DSMZ - Deutsche Sammlung von Microorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig DE, Germany.

The deposits DSM 13537, DSM 13538 and DSM 13539 were made by Novo Nordisk A/S and were later assigned to Novozymes A/S.

Polypeptide and polynucleotide sequences

The lipolytic enzyme of the invention may be a polypeptide having the amino acid sequence shown in SEQ ID NO: 2, 4, 6, 8 or 10 or it may be an amino acid variant derived from any of these sequences by substitution, deletion and/or insertion of one or more amino acid residues. The variant may be constructed in analogy with known variants of other lipases from *Fusarium*, e.g. from *F. oxysporum* and *F. heterosporum* as disclosed in WO 200032758 and T. Nagao, Journal of Bioscience and Bioengineering, 89 (5), 446-450 (2000).

The lipolytic enzyme may be C-terminally processed, e.g. by KEX-2 hydrolysis of the bond after Arg and Lys corresponding to K274 and R275 of SEQ ID NO: 6 to remove a peptide at the C-terminal. The C-terminally processed polypeptide tends to have a lower thermostability than the full-length polypeptide.

The variant may have substitutions of amino acids Arg and/or Lys corresponding to K274 and/or R275 of SEQ ID NO: 6, e.g. R275A and/or K274A. Such variants may be protected against C-terminal processing by KEX-2 hydrolysis to preserve the full-length polypeptide with higher thermostability.

A polynucleotide sequence encoding any of the polypeptides described above may be constructed by principles known in the art.

Properties of lipolytic enzyme

The lipolytic enzyme is able to hydrolyze carboxylic ester bonds and is classified as EC 3.1.1 according to Enzyme Nomenclature 1992, Academic Press, Inc. The enzyme has lipase (triacylglycerol lipase) activity (EC 3.1.1.3) and may also have other activities such as phospholipase activity, particularly phospholipase A1 (EC 3.1.1.32), and/or galactolipase (EC 3.1.1.26).

Recombinant expression vector

The expression vector of the invention typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal,

and, optionally, a selectable marker, a transcription terminator, a repressor gene or various activator genes. The vector may be an autonomously replicating vector, or it may be integrated into the host cell genome.

Production by cultivation of transformant

- 5 The lipolytic enzyme of the invention may be produced by transforming a suitable host cell with a DNA sequence encoding the lipolytic enzyme, cultivating the transformed organism under conditions permitting the production of the enzyme, and recovering the enzyme from the culture.

The host organism may particularly be a eukaryotic cell, in particular a fungal
10 cell, such as a yeast cell or a filamentous fungal cell, e.g. a strain of *Aspergillus*, *Fusarium*, *Trichoderma* or *Saccharomyces*, particularly *A. niger*, *A. oryzae*, *F. graminearum*, *F. sambucinum*, *F. cerealis* or *S. cerevisiae*. The production of the lipolytic enzyme in such host organisms may be done by the general methods described in EP 238,023 (Novo Nordisk), WO 96/00787 (Novo Nordisk) or EP
15 244,234 (Alko).

DNA recombination (shuffling)

The invention provides production of a lipolytic enzyme by shuffling of at least two polynucleotides, expressing the shuffled polynucleotides to form recombinant polypeptides, screening the polypeptides to select a polypeptide having
20 lipolytic enzyme activity, and producing the selected polypeptide.

The polynucleotides to be shuffled include a first polynucleotide comprising the and a second different polynucleotide encoding a lipolytic enzyme and having at least 50 % identity to the first polynucleotide. The second polynucleotide may comprise a second mature polypeptide-coding sequence of SEQ ID NO: 1, 3, 5, 7 or
25 9 or may comprise a polynucleotide encoding a lipolytic enzyme from *F. oxysporum* (WO 98/26057) or from *F. heterosporum*.

Shuffling (or recombination) of the homologous input polynucleotides (starting-point polynucleotides) may involve fragmenting the polynucleotides and recombining the fragments, to obtain output polynucleotides (i.e. polynucleotides
30 that have been subjected to a shuffling cycle) wherein a number of nucleotide fragments are exchanged in comparison to the input polynucleotides.

DNA recombination or shuffling may be a (partially) random process in which a library of chimeric genes is generated from two or more starting genes. A number of known formats can be used to carry out this shuffling or recombination process.

The process may involve random fragmentation of parental DNA followed by reassembly by PCR to new full length genes, e.g. as presented in US5605793, US5811238, US5830721, US6117679. In-vitro recombination of genes may be carried out, e.g. as described in US6159687, WO98/41623, US6159688, 5 US5965408, US6153510. The recombination process may take place *in vivo* in a living cell, e.g. as described in WO 97/07205 and WO 98/28416.

The parental DNA may be fragmented by DNA'se I treatment or by restriction endonuclease digests as described by Kikuchi et al (Gene 236:159-167, 2000). Shuffling of two parents may be done by shuffling single stranded parental DNA of 10 the two parents as described in Kikuchi et al (Gene 243:133-137, 2000).

A particular method of shuffling is to follow the methods described in Crameri et al, 1998, Nature, 391: 288-291 and Ness et al. Nature Biotechnology 17: 893-896. Another format would be the methods described in US 6159687: example 1 and 2.

Hybridization

15 The hybridization is used to indicate that a given DNA sequence is analogous to a nucleotide probe corresponding to a DNA sequence of the invention. The hybridization conditions are described in detail below.

Suitable conditions for determining hybridization between a nucleotide probe and a homologous DNA or RNA sequence involves presoaking of the filter 20 containing the DNA fragments or RNA to hybridize in 5 x SSC (standard saline citrate) for 10 min, and prehybridization of the filter in a solution of 5 x SSC (Sambrook et al. 1989), 5 x Denhardt's solution (Sambrook et al. 1989), 0.5 % SDS and 100 µg/ml of denatured sonicated salmon sperm DNA (Sambrook et al. 1989), followed by hybridization in the same solution containing a random-primed 25 (Feinberg, A. P. and Vogelstein, B. (1983) *Anal. Biochem.* 132:6-13), ³²P-dCTP-labeled (specific activity > 1 x 10⁹ cpm/µg) probe for 12 hours at approx. 45°C. The filter is then washed two times for 30 minutes in 2 x SSC, 0.5 % SDS at a temperature of at least 55°C, more particularly at least 60°C, more particularly at least 65°C, even more particularly at least 70°C, especially at least 75°C.

30 Molecules to which the oligonucleotide probe hybridizes under these conditions are detected using a x-ray film.

Alignment and identity

The lipolytic enzyme and the nucleotide sequence of the invention may have homologies to the disclosed sequences of at least 85 %, particularly at least 90 % or 35 at least 95 %, e.g. at least 98 %.

For purposes of the present invention, alignments of sequences and calculation of identity scores were done using a Needleman-Wunsch alignment (i.e. global alignment), useful for both protein and DNA alignments. The default scoring matrices BLOSUM50 and the identity matrix are used for protein and DNA alignments respectively. The penalty for the first residue in a gap is -12 for proteins and -16 for DNA, while the penalty for additional residues in a gap is -2 for proteins and -4 for DNA. Alignment is from the FASTA package version v20u6 (W. R. Pearson and D. J. Lipman (1988), "Improved Tools for Biological Sequence Analysis", PNAS 85:2444-2448, and W. R. Pearson (1990) "Rapid and Sensitive Sequence Comparison with FASTP and FASTA", Methods in Enzymology, 183:63-98).

Lipase activity (LU)

A substrate for lipase is prepared an emulsion of 5 % by volume of tributyrin (glycerin tributyrate) using 0.1 % gum Arabic as emulsifier. The hydrolysis of tributyrin at 30 °C at pH 7 is followed in a pH-stat titration experiment. One unit of lipase activity (1 LU) equals the amount of enzyme capable of releasing 1 μ mol butyric acid/min at the standard conditions. 1 KLU = 1000 LU.

Use of lipolytic enzyme

The lipolytic enzyme of the invention can be used in various industrial application of lipolytic enzymes, e.g. in baking or detergents as described below. It can also be used for diglyceride synthesis (EP 307154), acidolysis, interesterification (WO 8802775), ester hydrolysis, oil degumming (JP-A 2-153997, US 5264367), production of lysolecithin (JP patent 2794574, JP-B 6-087751) and in a cheese-making process (WO 0054601).

25 Use in baking

The lipolytic enzyme of the invention can be used in the preparation of dough, bread and cakes, e.g. to improve the elasticity of the bread or cake. Thus, the lipolytic enzyme can be used in a process for making bread, comprising adding the lipolytic enzyme to the ingredients of a dough, kneading the dough and baking the dough to make the bread. This can be done in analogy with WO 9404035 and EP 585988. The lipolytic enzyme is typically added in an amount corresponding to 0.01-100 mg enzyme protein per kg of flour, particularly 0.1-25 mg enzyme protein per kg of flour, more particularly 0.1-10 mg enzyme protein per kg of flour, and most particularly 0.5-5 mg enzyme protein per kg of flour.

Use in detergent

The lipolytic enzyme may be used as a detergent additive, e.g. at a concentration (expressed as pure enzyme protein) of 0.001-10 (e.g. 0.01-1) mg per gram of detergent or 0.001-100 (e.g. 0.01-10) mg per liter of wash liquor.

5 The detergent composition of the invention may for example be formulated as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations. In a laundry detergent, the lipolytic
10 enzyme may be effective for the removal of fatty stains, for whiteness maintenance and for dingy cleanup. A laundry detergent composition may be formulated as described in WO 97/04079, WO 97/07202, WO 97/41212, PCT/DK WO 98/08939 and WO 97/43375.

The detergent composition of the invention may particularly be formulated
15 for hand or machine dishwashing operations. e.g. as described in GB 2,247,025 (Unilever) or WO 99/01531 (Procter & Gamble). In a dishwashing composition, the lipolytic enzyme may be effective for removal of greasy/oily stains, for prevention of the staining /discoloration of the dishware and plastic components of the dishwasher by highly colored components and the avoidance of lime soap deposits on the
20 dishware.

MATERIALS AND METHODS

Methods

Unless otherwise stated, DNA manipulations and transformations were performed using standard methods of molecular biology as described in Sambrook
25 et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology", John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and Sons, 1990.

Enzymes

30 Enzymes for DNA manipulations (e.g. restriction endonucleases, ligases etc.) are obtainable from New England Biolabs, Inc. and were used according to the manufacturer's instructions.

Plasmids/vectors

pT7Blue (Invitrogen, Netherlands)

pCaHj483 is described in WO 9704079 and WO 9942566.

Cloning

LA PCR™ in vitro Cloning Kit (TaKaRa) was used for cloning and was used according to the manufacturer's instructions.

5 Microbial strains

Fusarium venenatum WTY700 3.8d, a spore-purified *tri5*-minus, *dps1*-minus strain, was used as the recipient strain for transformation experiments. *Fusarium venenatum* WTY700 3.8d is a morphological mutant of *Fusarium venenatum* strain ATCC 20334 (Wiebe *et al.*, 1991, *Mycol. Research* 95: 1284-1288),

10 *E. coli* JM109 (TOYOBO, Japan)

E. coli JM110 (Invitrogen)

A. oryzae BECh-2 is described in Danish patent application PA 1999 01726. It is a mutant of JaL 228 (described in WO 98/12300) which is a mutant of IFO 4177.

Media and reagents

15 Cove: 342.3 g/L Sucrose, 20 ml/L COVE salt solution, 10mM Acetamide, 30 g/L noble agar.

Cove-2: 30 g/L Sucrose, 20 ml/L COVE salt solution, 10mM, Acetamide, 30 g/L noble agar.

Cove salt solution: per liter 26 g KCl, 26 g MgSO4-7aq, 76 g KH2PO4, 50ml
20 Cove trace metals.

Cove trace metals: per liter 0.04 g NaB4O7-10aq, 0.4 g CuSO4-5aq, 1.2 g FeSO4-7aq, 0.7 g MnSO4-aq, 0.7 g Na2MoO2-2aq, 0.7 g ZnSO4-7aq.

AMG trace metals: per liter 14.3 g ZnSO4-7aq, 2.5 g CuSO4-5aq, 0.5 g NiCl2, 13.8 g FeSO4, 8.5 g MnSO4, 3.0 g citric acid.

25 YPG: 4 g/L Yeast extract, 1 g/L KH2PO4, 0.5 g/L MgSO4-7aq, 5 g/L Glucose, pH 6.0.

STC: 0.8 M Sorbitol, 25 mM Tris pH 8, 25 mM CaCl2.

STPC: 40 % PEG4000 in STC buffer.

Cove top agarose: 342.3 g/L Sucrose, 20 ml/L COVE salt solution, 10mM
30 Acetamide, 10 g/L low melt agarose.

MS-9: per liter 30 g soybean powder, 20 g glycerol, pH 6.0.

MDU-pH5: per liter 45 g maltose-1aq, 7 g yeast extract, 12 g KH2PO4, 1 g MgSO4-7aq, 2 g K2SO4, 0.5 ml AMG trace metal solution and 25 g 2-morpholinoethanesulfonic acid, pH 5.0.

Measurement of dough and breads properties

The crumb firmness was measured using a texture analyzer TA-XT2 from Stable Micro Systems.

Texture was measured according to a modified ACCA method (American
5 Cereal Chemists' Association).

Elasticity is a measure of the degree to which a dough can be stretched without tearing. It is evaluated by rolling a piece of dough (about 30 g) to a size of about 10 cm, and having a test panel carefully pulling at opposite ends to judge the resistance and elasticity. The results are expressed on a scale from 0 (low/weak
10 elasticity) to 10 (high/strong elasticity) with the control (dough without enzyme addition) taken as 5.

Crumb structure is determined as the property of a baked product with finer and/or thinner cell walls in the crumb and/or more uniform/homogenous distribution of cells in the crumb and is usually evaluated empirically by the skilled test baker.

15 The shape factor is taken as the ratio between the height and diameter of rolls after baking (average of 10 rolls).

The specific volume is calculated as volume ml per g bread, taking the mean value of the volumes of 4 loaves measured using the traditional rape seed method. The specific volume of the control (without enzyme) is defined as 100. The relative
20 specific volume index is the percentage of the specific volume of 4 loaves per the specific volume of 4 control loaves.

EXAMPLES

Example 1: Fermentation and mycelial tissue from *F. venenatum*

Fusarium venenatum WTY700 3.8d was grown in a two-liter lab-scale
25 fermentor using a fed-batch fermentation scheme with NUTRIOSE™ (Roquette Freres, S.A., Beinheim, France) as the carbon source and yeast extract. Ammonium phosphate was provided in the feed. The fermentation was maintained at pH 6-6.5 and 30°C with positive dissolved oxygen.

Mycelial samples were harvested at 2, 4, 6, and 8 days post-inoculum and
30 quick-frozen in liquid nitrogen. The samples were stored at -80°C until they were disrupted for RNA extraction.

Example 2: cDNA library construction

Total cellular RNA was extracted from the mycelial samples described in Example 1 according to the method of Timberlake and Barnard (1981, *Cell* 26: 29-

37), and the RNA samples were analyzed by Northern hybridization after blotting from 1% formaldehyde-agarose gels (Davis *et al.*, 1986, *Basic Methods in Molecular Biology*, Elsevier Science Publishing Co., Inc., New York). Polyadenylated mRNA fractions were isolated from total RNA with an mRNA Separator Kit™ (Clontech
5 Laboratories, Inc., Palo Alto, CA) according to the manufacturer's instructions. Double-stranded cDNA was synthesized using approximately 5 µg of poly(A)+ mRNA according to the method of Gubler and Hoffman (1983, *Gene* 25: 263-269) except a *NotI*-(dT)18 primer (Pharmacia Biotech, Inc., Piscataway, NJ) was used to initiate first strand synthesis. The cDNA was treated with mung bean nuclease
10 (Boehringer Mannheim Corporation, Indianapolis, IN) and the ends were made blunt with T4 DNA polymerase (New England Biolabs, Beverly, MA).

The cDNA was digested with *NotI*, size selected by agarose gel electrophoresis (ca. 0.7-4.5 kb), and ligated with pZER0-2.1 (Invitrogen Corporation, Carlsbad, CA) which had been cleaved with *NotI* plus *EcoRV* and dephosphorylated
15 with calf-intestine alkaline phosphatase (Boehringer Mannheim Corporation, Indianapolis, IN). The ligation mixture was used to transform competent *E. coli* TOP10 cells (Invitrogen Corporation, Carlsbad, CA). Transformants were selected on 2YT agar plates (Miller, 1992, *A Short Course in Bacterial Genetics. A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria*, Cold Spring Harbor
20 Press, Cold Spring Harbor, New York) which contained kanamycin at a final concentration of 50 µg/ml.

Example 3: Template preparation and nucleotide sequencing

From the cDNA library described in Example 2, 1192 transformant colonies were picked directly from the transformation plates into 96-well microtiter dishes
25 which contained 200 µl of 2YT broth (Miller, 1992, *supra*) with 50 µg/ml kanamycin. The plates were incubated overnight at 37°C without shaking. After incubation 100 µl of sterile 50% glycerol was added to each well. The transformants were replicated into secondary, deep-dish 96-well microculture plates (Advanced Genetic Technologies Corporation, Gaithersburg, MD) containing 1 ml of Magnificent Broth™
30 (MacConnell Research, San Diego, CA) supplemented with 50 µg of kanamycin per ml in each well. The primary microtiter plates were stored frozen at -80°C. The secondary deep-dish plates were incubated at 37°C overnight with vigorous agitation (300 rpm) on rotary shaker. To prevent spilling and cross-contamination, and to allow sufficient aeration, each secondary culture plate was covered with a
35 polypropylene pad (Advanced Genetic Technologies Corporation, Gaithersburg, MD) and a plastic microtiter dish cover.

DNA was isolated from each well using the 96-well Miniprep Kit protocol of Advanced Genetic Technologies Corporation (Gaithersburg, MD) as modified by Utterback *et al.* (1995, *Genome Sci. Technol.* 1: 1-8). Single-pass DNA sequencing (EST) was done with a Perkin-Elmer Applied Biosystems Model 377 XL Automated DNA Sequencer (Perkin-Elmer/Applied Biosystems, Inc., Foster City, CA) using dye-terminator chemistry (Giesecke *et al.*, 1992, *Journal of Virology Methods* 38: 47-60) and the reverse lac sequencing primer.

Example 4: Analysis of DNA sequence data

Nucleotide sequence data were scrutinized for quality, and samples giving improper spacing or ambiguity levels exceeding 3% were discarded or re-run. Vector sequences and ambiguous base calls at the ends of the DNA sequences were trimmed with assistance of FACTURA™ software (Perkin-Elmer Applied Biosystems, Inc., Foster City, CA). All sequences were compared to each other to determine multiplicity using AutoAssembler™ software (Perkin-Elmer Applied Biosystems, Inc., Foster City, CA). Lastly, all sequences were translated in three frames and searched against a non-redundant database (NRDB) using GeneAssist™ software (Perkin-Elmer Applied Biosystems, Inc., Foster City, CA) with a modified Smith-Waterman algorithm using the BLOSUM 62 matrix with a threshold score of 70. The NRDB was assembled from Genpept, Swiss-Prot, and PIR databases.

Example 5: Identification of lipase cDNA clones

Putative lipase clones were identified by comparing the deduced amino acid sequence of the ESTs to protein sequences deposited in publicly available databases such as Swissprot, Genpept, and PIR using a modified Smith-Waterman search algorithm (Perkin-Elmer Applied Biosystems, Foster City, CA). Tentative identification was based on amino acid sequence similarity to numerous fungal lipases. One clone, *Fusarium venenatum* EST FA0726, was selected for nucleotide sequence analysis which revealed that the cDNA clone was truncated at its 5 prime end.

Example 6: *Fusarium venenatum* genomic DNA extraction

Fusarium venenatum WTY700 was grown for 24 hours at 28°C and 150 rpm in 25 ml of YEG medium composed per liter of 5 g of yeast extract and 20 g of glucose. Mycelia were then collected by filtration through Miracloth (Calbiochem, La Jolla, CA) and washed once with 25 ml of 10 mM Tris-1 mM EDTA (TE) buffer. Excess buffer was drained from the mycelia which were subsequently frozen in liquid

nitrogen. The frozen mycelia were ground to a fine powder in an electric coffee grinder, and the powder was added to 20 ml of TE buffer and 5 ml of 20% w/v sodium dodecylsulfate (SDS) in a disposable plastic centrifuge tube. The mixture was gently inverted several times to ensure mixing, and extracted twice with an equal volume of phenol:chloroform:isoamyl alcohol (25:24:1 v/v/v). Sodium acetate (3 M solution) was added to give a final concentration of 0.3 M and the nucleic acids were precipitated with 2.5 volumes of ice cold ethanol. The tube was centrifuged at 15,000 x g for 30 minutes and the pellet was allowed to air dry for 30 minutes before resuspension in 0.5 ml of TE buffer. DNase-free ribonuclease A was added to a concentration of 100 µg/ml and the mixture was incubated at 37°C for 30 minutes. Proteinase K (200 µg/ml) was then added and the mixture was incubated an additional hour at 37°C. Finally, the mixture was extracted twice with phenol:chloroform:isoamyl alcohol (25:24:1 v/v/v) before precipitating the DNA with sodium acetate and ethanol according to standard procedures. The DNA pellet was dried under vacuum, resuspended in TE buffer, and stored at 4°C.

Example 7: Genomic DNA library construction, screening, and isolation of genomic Lipase clone

Genomic libraries of *Fusarium venenatum* WTY700 were constructed in λZipLox according to the manufacturer's instructions (Life Technologies, Gaithersburg, MD). *Fusarium venenatum* genomic DNA was partially digested with *Tsp509I* and size-fractionated on 1% agarose gels. DNA fragments migrating in the size range 3-7 kb were excised and eluted from the agarose gel slices using Prep-a-Gene reagents (BioRad, Hercules, CA). The eluted DNA fragments were ligated with *EcoRI*-cleaved and dephosphorylated λZipLox vector arms (Life Technologies, Gaithersburg, MD), and the ligation mixtures were packaged using commercial packaging extracts (Stratagene, La Jolla, CA). The packaged DNA libraries were plated and amplified in *E. coli* Y1090ZL cells.

The cDNA from *Fusarium venenatum* clone FA0726 was excised from the vector plasmid by digestion with *EcoRI* and *NotI* yielding an approximately 900 bp fragment. The fragment was purified by gel electrophoresis, and radiolabeled with α[³²P] dCTP using a Prime-it Random Primer Labeling Kit (Stratagene, La Jolla, CA).

Approximately 40,000 plaques from the library were screened by plaque-hybridization (Davis *et al.*, 1980, *supra*) with the radiolabeled probe fragment of the *Fusarium venenatum* lipase gene using high stringency conditions at 45°C (high stringency = 50% formamide, 5X SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA). Plaques, which gave hybridization signals, were

purified once in *E. coli* DH10B cells, and the individual clones were subsequently excised from the λ ZipLox vector as pZL1-derivatives (D'Alessio *et al.*, 1992, *Focus*® 14: 7).

One plaque was identified that hybridized strongly to the *Fusarium*
5 *venenatum* lipase gene probe, and was subsequently excised from the λ ZipLox vector as a pZL1-derivative (D'Alessio *et al.*, 1992, *supra*). Plasmid DNA was isolated from the clone by passage through *E. coli* DH10B cells (Life Technologies, Gaithersburg, MD) according to the manufacturer's instructions. This clone was designated *E. coli* DH10B – pFvLipase1.

10 **Example 8: Characterization of the *Fusarium venenatum* genomic clone encoding lipase**

DNA sequencing was performed on an Perkin-Elmer Biosystems Model 377 XL Automated DNA Sequencer using dye-terminator chemistry (Giesecke *et al.*, 1992, *Journal of Virology Methods* 38: 47-60). Contig sequences were generated using a
15 transposon insertion strategy (Primer Island Transposition Kit, Perkin-Elmer/Applied Biosystems, Inc., Foster City, CA). The 2.94 kb genomic fragment was sequenced to an average redundancy of 4.8.

The nucleotide sequence and deduced amino acid sequence are shown as SEQ ID NO: 1-2. The insert contains an open reading frame of 1.153 kb encoding a
20 polypeptide of 350 amino acids. Using the SignalP software program (Nielsen *et al.*, 1997, *Protein Engineering* 10: 1-6), a signal peptide of 15 residues was predicted. The predicted signal peptide is followed by a 15 residue propeptide ending with a concanonical propeptide Glu/Arg cleavage site. N-terminal sequencing of the lipase protein supports this propeptide cleavage site prediction. The open reading frame is
25 interrupted by two introns of 49 bp and 58 bp. Thus, the mature *Fusarium venenatum* lipase comprises 319 amino acids and a predicted molecular weight of 33.6 kDa. There are 2 potential N-linked glycosylation sites (Asn-X-Ser/Thr) within the *Fusarium venenatum* lipase.

A comparative alignment of lipase sequences using the Clustal W algorithm
30 in the Megalign program of DNA-Star, showed that the deduced amino acid sequence of the *Fusarium venenatum* lipase gene shares 81% identity to the deduced amino acid sequence of a *Fusarium oxysporum* phospholipase A (EP0869167).

Example 9: Construction of plasmid pSheB1

The *Fusarium venenatum* expression vector pSheB1 (Figure 1) was generated by modification of pDM181 (WO 98/20136). The modifications included (a) removal of two *Nco*I sites within the pDM181 sequence, and (b) restoration of the natural translation start of the *Fusarium oxysporum* trypsin promoter (reconstruction
5 of an *Nco*I site at the ATG start codon).

Removal of two *Nco*I sites within the pDM181 sequence was accomplished using the QuikChange™ site-directed mutagenesis kit (Stratagene Cloning Systems, La Jolla, CA) according to the manufacturer's instruction with the following pairs of mutagenesis primers: SEQ ID NO: 11 and 12, SEQ ID NO: 13 and 14.

10 Restoration of the natural translation start of the *Fusarium oxysporum* trypsin promoter was also accomplished using the Stratagene QuikChange™ site directed mutagenesis kit in conjunction with the following pair of mutagenesis primers: SEQ ID NO:15 and 16.

All site-directed changes were confirmed by DNA sequence analysis of the
15 appropriate vector regions.

Example 10: Construction of expression vector pEJG60

The lipase-expression vector, pEJG60 was constructed as follows. The lipase coding region was amplified from pFvlipase1 using the following pair of primers: Primer 990658 and 990661 (SEQ ID NO: 17 and 18)

20 The forward primer introduces a *Sph*I site which contains the ATG, and the reverse primer introduces a *Pac*I site after the stop codon.

The amplification reaction (100 µl) contained the following components: 0.5 µg of genomic clone pFvLipase1, 50 pmol of the forward primer, 50 pmol of the reverse primer, 10 mM dNTPs (dATP, dCTP, dGTP, and dTTP), 1X *Pwo* DNA
25 polymerase buffer, and 2.5 units of *Pwo* DNA polymerase (Boehringer Mannheim, Indianapolis, IN). The reactions were incubated in a Perkin-Elmer Model 480 Thermal Cycler programmed for 1 cycles at 95°C for 2 minutes; 10 cycles at 94°C for 45 seconds, 55°C for 45 seconds, and 72°C for 2 minutes; 17 cycles at 94°C for 45 seconds, 55°C for 45 seconds, and 72°C for 2 minutes with an extension of 20
30 seconds per cycle; 1 cycle at 72°C for 10 minutes; and a soak cycle at 4 °C. The reaction products were isolated on a 1% agarose gel where a 1.15 kb product band was excised from the gel and purified using Qiaquick Gel Extraction Kit (Qiagen, Chatsworth, CA) according to the manufacturer's instructions.

The generated fragment was digested with *Sph*I, blunted with Klenow,
35 digested with *Pac*I, and purified by agarose gel electrophoresis and Qiaquick Gel Extraction Kit (Qiagen, Chatsworth, CA). The purified DNA segment was ligated into

pSheB1 (Figure 1) which was previously *Nco*I digested, treated with DNA polymerase I (Klenow fragment), and digested with *Pac*I. The treatment of the *Nco*I-digested vector with Klenow fragment resulted in a filling in of the *Nco*I cohesive end, thereby making it blunt and compatible with the blunt site of the lipase DNA segment. The resulting expression plasmid was designated pEJG60 (Figure 2). The PCR-amplified lipase gene segment was re-sequenced to verify the absence of any errors.

Example 11: Transformation of *Fusarium venenatum* and analysis of *Fusarium venenatum* transformants

10 Spores of *Fusarium venenatum* WTY700 were generated by inoculating a flask containing 500 ml of RA sporulation medium with 10 plugs from a 1X Vogels medium plate (2.5% Noble agar) supplemented with 2.5% glucose and 2.5 mM sodium nitrate and incubating at 28°C, 150 rpm for 2 to 3 days. Spores were harvested through Miracloth (Calbiochem, San Diego, CA) and centrifuged 20
15 minutes at 7000 rpm in a Sorvall RC-5B centrifuge (E. I. DuPont De Nemours and Co., Wilmington, DE). Pelleted spores were washed twice with sterile distilled water, resuspended in a small volume of water, and then counted using a hemocytometer.

Protoplasts were prepared by inoculating 100 ml of YEPG medium with 4×10^7 spores of *Fusarium venenatum* WTY700 and incubating for 16 hours at 24°C
20 and 150 rpm. The culture was centrifuged for 7 minutes at 3500 rpm in a Sorvall RT 6000D (E. I. DuPont De Nemours and Co., Wilmington, DE). Pellets were washed twice with 30 ml of 1 M MgSO_4 and resuspended in 15 ml of 5 mg/ml of NOVOZYME 234™ (batch PPM 4356, Novo Nordisk A/S, Bagsværd, Denmark) in 1 M MgSO_4 . Cultures were incubated at 24°C and 150 rpm until protoplasts formed. A volume of
25 35 ml of 2 M sorbitol was added to the protoplast digest and the mixture was centrifuged at 2500 rpm for 10 minutes. The pellet was resuspended, washed twice with STC, and centrifuged at 2000 rpm for 10 minutes to pellet the protoplasts. Protoplasts were counted with a hemocytometer and resuspended in an 8:2:0.1 solution of STC:SPTC:DMSO to a final concentration of 1.25×10^7 protoplasts/ml.
30 The protoplasts were stored at -80°C, after controlled-rate freezing in a Nalgene Cryo 1°C Freezing Container (VWR Scientific, Inc., San Francisco, CA).

Frozen protoplasts of *Fusarium venenatum* WTY700 were thawed on ice. Five μg of pEJG60 described in Example 10 and 5 μl of heparin (5 mg per ml of STC) was added to a 50 ml sterile polypropylene tube. One hundred μl of
35 protoplasts was added, mixed gently, and incubated on ice for 30 minutes. One ml of SPTC was added and incubated 20 minutes at room temperature. After the addition

of 25 ml of 40°C COVE top agarose, the mixture was poured onto an empty 150 mm diameter plate and incubated overnight at room temperature. Then an additional 25 ml of 40°C COVE top agarose containing 10 mg of BASTA™ per ml was poured on top of the plate and incubated at room temperature for up to 14 days. The active
5 ingredient in the herbicide BASTA™ is phosphinothricin. BASTA™ was obtained from AgrEvo (Hoechst Schering, Rodovre, Denmark) and was extracted twice with phenol:chloroform:isoamyl alcohol (25:24:1), and once with chloroform:isoamyl alcohol (24:1) before use.

Twenty-four transformants were picked directly from the selection plates
10 (COVE underlay with COVE-BASTA™ overlay) and inoculated into 125 ml shake flasks containing 25 ml of M400Da medium supplemented with 1 mM CaCl₂ and 100 µg/ml ampicillin (to prevent bacterial contamination) and incubated at 28°C, 200 rpm on a platform shaker for 7 days. The untransformed recipient strain was also included as a negative control.

15 Flasks were sampled at 5 and 7 days and assayed for lipase activity as described below. The samples were also submitted to SDS-PAGE using Novex gradient gels (Novex Experimental Technology, San Diego, CA).

Lipase activity was determined as follows: 100 µl of substrate (3.92 ml of 100 mM MOPS pH 7.5, 4 mM CaCl₂, 990 µl of DMSO, 80 µl of 1% AOS, and 20 µl of p-
20 nitrophenyl butyrate) was added to 100 µl of diluted sample. The samples were diluted accordingly in 100 mM MOPS pH 7.5, 4 mM CaCl₂. The absorbance at 405 nm was monitored for 3 minutes at room temperature in a 96-well microtiter plate using a Molecular Devices Thermomax Microplate Reader.

The lipase assay results indicated that at both 5 and 7 days, most of the
25 transformants produced lipase activity well above that of the untransformed control. Shake flask culture broths from transformants #1 and #3, the two highest scorers in the lipase assay, were analyzed on a 16% tricine gel. A prominent polypeptide at a apparent molecular weight of 32-33 kD was observed at both time points and for each transformant harboring pEJG60.

30 **Example 12: Cloning and expression of lipase gene from *Fusarium sulphureum***

Transformation in *Aspergillus* strain

Aspergillus oryzae strain BECh-2 was inoculated to 100 ml of YPG medium and incubated for 16 hrs at 32°C at 120 rpm. Pellets were collected and washed with
35 0.6 M KCl, and resuspended in 20 ml 0.6 M KCl containing a commercial β-glucanase product (Glucanex, product of Novo Nordisk A/S) at the concentration of 30 µl/ml.

Cultures were incubated at 32°C at 60 rpm until protoplasts formed, then washed with STC buffer twice. The protoplasts were counted with a hematometer and resuspended in an 8:2:0.1 solution of STC:STPC:DMSO to a final concentration of 2.5x10⁷ protoplasts/ml. About 3 µg of DNA was added to 100 µl of protoplasts solution, mixed gently and incubated on ice for 30 min. One ml of SPTC was added and incubated 30 min at 37°C. After the addition of 10 ml of 50°C Cove top agarose, the reaction was poured onto Cove agar plate. Transformation plates were incubated at 32°C for 5 days.

PCR screening of lipase

10 A strain of *Fusarium sulphureum* was used as a genomic DNA supplier.

PCR reactions on *Fusarium sulphureum* genomic DNA was done with two following primer sets: lip3 / lip15 (SEQ ID NO: 20/23) and lip10 / lip17 (SEQ ID NO: 21/24) designed based upon the alignment of 3 lipases from *Fusarium*.

Reaction components (2.6 ng /µl of genomic DNA, 250 mM dNTP each, 15 primer 250 nM each, 0.1 U/ µl of Taq polymerase in 1X buffer (Roche Diagnostics, Japan)) were mixed and submitted for PCR under the following conditions.

Step	Temperature	Time
1	94°C	1 min
2	50°C	1 min
3	72°C	2 min
4	72°C	10 min
5	4°C	forever

Steps 1 to 3 were repeated 30 times.

450bp of fragment and 280 bp of fragment were amplified from primer sets of lip3/lip15 (SEQ ID NO: 20/23) and lip10/lip17 (SEQ ID NO: 21/24), respectively. They were gel-purified with GFXTM PCR DNA and Gel Band Purification kit (amersham pharmacia biotech) and ligated into a pT7Blue vector with ligation high (TOYOBO, Japan). The ligation mixtures were transformed into *E. coli* JM109. The resultant plasmids, pT27-0315 and pT27-1017, were sequenced and compared to the *Fusarium oxysporum* lipase, showing that a clone encodes the internal part of the lipase.

Cloning of lipase gene

In order to clone the missing part of the lipase gene, adaptor PCR was done. A cassette was made by mixing of adaptor L (SEQ ID NO: 26) and adaptor S:: acctgccc.

- 5 3' and 5' of adaptor S are dephosphorylated and amidated, respectively.

1.3 µg of Eco RV digested genome was ligated to the cassette and it was used as a PCR template. Reaction components (7 ng /µl of genomic DNA ligated to cassette, 250 mM dNTP each, primer 250 nM each, 0.05 U/ µl of Expand high fidelity polymerase in 1X buffer (Roche Diagnostics, Japan)) were mixed and submitted for

- 10 PCR under the following conditions.

Step	Temperature	Time
1	94°C	2 min
2	94°C	10sec
3	55°C	30sec
4	68°C	45sec
step 2-4 repeat 10 times		
5	94°C	10sec
6	55°C	30sec
7	68°C	45sec +20sec/cycle
step 5-7, repeat 20 times		
8	68°C	7min
7	4°C	forever

- 500 bp of DNA fragment corresponding to N-terminal region was obtained with 27N1long primer (SEQ ID NO: 27) and 200 bp of DNA fragment corresponding to C-terminal region was obtained with 27C1long primer (SEQ ID NO: 28).

Obtained fragments were purified by GFXTM PCR DNA and Gel Band Purification kit (Amersham Pharmacia Biotech) and sequenced with each primers which amplified the fragment.

- The missing C-terminal part was cloned with LA PCRTM in vitro Cloning Kit (TaKaRa) following to the manufacturer's instructions. 350 bp of DNA fragment corresponding to C-terminal region was obtained from Xho I digested genome ligated to Sal I cassette of the kit with 27C2 primer (SEQ ID NO: 29).

Obtained fragments were purified by GFX™ PCR DNA and Gel Band Purification kit (Amersham Pharmacia Biotech) and sequenced with 27C2 primer.

The fidelity of taq polymerase is not good so in order to get the right sequence whole gene was amplified the following primers: 27N(Bam) and 27C(Sal)
5 (SEQ ID NO: 30 and 31).

Reaction components (6 ng / μ l of genomic DNA, 250 mM dNTP each, primer 250 nM each, 0.05 U/ μ l of Expand high fidelity polymerase in 1X buffer (Roche Diagnostics, Japan)) were mixed and submitted for PCR under the following conditions.

10

Step	Temperature	Time
1	94°C	2 min
2	94°C	10sec
3	55°C	30sec
4	68°C	45sec
step 2-4 repeat 10 times		
5	94°C	10sec
6	55°C	30sec
7	68°C	45sec +20sec/cycle
step 5-7, repeat 20 times		
8	68°C	7min
7	4°C	forever

Amplified DNA fragment was gel-purified with GFX™ PCR DNA and Gel Band Purification kit (amersham pharmacia biotech) and ligated into a pT7Blue vector with ligation high (TOYOBO, Japan). The ligation mixtures were transformed
15 into *E. coli* JM109. Four plasmids, pT27w-1, pT27w-2, pT27w-3, and pT27w-4, were sequenced and their sequence were compared. pT27w-3 has no PCR error and it is defined as *Fusarium sulphureum* lipase nucleotide sequence.

Expression of lipase gene in *Aspergillus oryzae*.

The lipase gene was digested from pT27w-3 with BamH I and Sal I and
20 ligated into the BamH I and XhoI sites in the *Aspergillus* expression cassette pCaHj483 which has *Aspergillus niger* neutral amylase promoter, *Aspergillus*

nidulans TPI leader sequences, *Aspergillus niger* glucoamylase terminator and *Aspergillus nidulans* amdS gene as a marker. The resultant plasmid was pNL27w-8.

pNL27w-8 was transformed into *Aspergillus oryzae* BECh-2. The selected transformants were inoculated in 100 ml of MS-9 media and cultivated at 30°C for 1 day. 3 ml of grown cell in MS-9 medium was inoculated to 100 ml of MDU-2BP medium and cultivated at 32°C for 3 days. The supernatant was obtained by centrifugation.

The lipase productivity of selected transformants was determined as LU activity. The productivity of the best transformant TNL27-75 was 130 LU/ml and BECh2 has no lipase activity.

Example 13: Immunological characterization of lipolytic enzyme from *Fusarium sulphureum*

A purified lipolytic enzyme sample having the amino acid sequence shown as amino acids 1-319 of SEQ ID NO: 4 was tested by immunodiffusion against a polyclonal antibody raised against the *Fusarium oxysporum* lipase. No immunological cross-reaction was found.

Example 14: Cloning and expression of lipase gene from *Acremonium berkeleyanum*

Transformation in *Aspergillus* strain

Aspergillus oryzae strain BECh-2 was inoculated to 100 ml of YPG medium and incubated for 16 hrs at 32°C at 120 rpm. Pellets were collected and washed with 0.6 M KCl, and resuspended in 20 ml 0.6 M KCl containing a commercial β -glucanase product (Glucanex, product of Novo Nordisk A/S) at the concentration of 30 μ l/ml. Cultures were incubated at 32°C at 60 rpm until protoplasts formed, then washed with STC buffer twice. The protoplasts were counted with a hematology counter and resuspended in an 8:2:0.1 solution of STC:STPC:DMSO to a final concentration of 2.5×10^7 protoplasts/ml. About 3 μ g of DNA was added to 100 μ l of protoplasts solution, mixed gently and incubated on ice for 30 min. One ml of STPC was added and incubated 30 min at 37°C. After the addition of 10 ml of 50°C Cope top agarose, the reaction was poured onto Cope agar plate. Transformation plates were incubated at 32°C for 5 days.

PCR screening of lipase

A strain of *Acremonium berkeleyanum* was used as a genomic DNA supplier.

PCR reactions on *Acremonium berkeleyanum* genomic DNA was done with two following primer sets: lip3 / lip11 (SEQ ID NO: 20/22) and lip10 / lip21 (SEQ ID NO: 21/25) designed based upon the alignment 3 lipases from *Fusarium*.

Reaction components (2.5 ng / μ l of genomic DNA, 250 mM dNTP each, 5 primer 250 nM each, 0.1 U/ μ l in Taq polymerase in 1X buffer (Roche Diagnostics, Japan)) were mixed and submitted for PCR under the following conditions.

Step	Temperature	Time
1	94°C	1 min
2	50°C	1 min
3	72°C	2 min
4	72°C	10 min
5	4°C	forever

Steps 1 to 3 were repeated 30 times.

340 bp of fragment and 330 bp of fragment were amplified from primer sets 10 of lip3/lip11 and lip10/lip21, respectively. They were gel-purified with GFX™ PCR DNA and Gel Band Purification kit (amersham pharmacia biotech) and ligated into a pT7Blue vector with ligation high (TOYOBO, Japan). The ligation mixtures were transformed into *E. coli* JM109. The resultant plasmids, pTAc-0310 and pTAc-1021, were sequenced and compared to the *Fusarium oxysporum* lipase, showing that a 15 clone encodes the internal part of the lipase.

Cloning of lipase gene

In order to clone the missing part of the lipase gene, LA PCR™ in vitro Cloning Kit (TaKaRa) was used for genome walking. One kb of DNA fragment corresponding to N-terminal region was obtained from Sal I digested genome ligated 20 to Sal I cassette of the kit with AcN3 primer (SEQ ID NO: 32). One kb of DNA fragment corresponding to C-terminal region was obtained from Hind III digested genome ligated to Hind III cassette of the kit with AcC3 primer (SEQ ID NO: 33).

Obtained fragments were purified by GFX™ PCR DNA and Gel Band Purification kit (amersham pharmacia biotech) and sequenced with each primer 25 which amplified the fragment. The sequences were compared to *Fusarium oxysporum* lipase, showing that an amplified DNA fragments encodes the whole part of lipase.

The fidelity of taq polymerase is not good so in order to get the right sequence whole gene was amplified the primers 152-N (Bcl) and 152-C(Xho) (SEQ ID NO: 34 and 35).

Reaction components (6 ng / μ l of genomic DNA, 250 mM dNTP each, primer 250 nM each, 0.05 U/ μ l of Expand high fidelity polymerase in 1X buffer (Roche Diagnostics, Japan)) were mixed and submitted for PCR under the following conditions.

5

Step	Temperature	Time
1	94°C	2 min
2	94°C	10sec
3	55°C	30sec
4	68°C	45sec
step 2-4 repeat 10 times		
5	94°C	10sec
6	55°C	30sec
7	68°C	45sec +20sec/cycle
step 5-7, repeat 20 times		
8	68°C	7min
7	4°C	forever

- 10 Amplified DNA fragment was gel-purified with GFX™ PCR DNA and Gel Band Purification kit (amersham pharmacia biotech) and ligated into a pT7Blue vector with ligation high (TOYOBO, Japan). The ligation mixtures were transformed into *E. coli* JM109. The resultant plasmids, pT152-1 and pT152-2, were sequenced and they are identical. pT152-1 sequence is defined as *Acremonium berkeleyanum* lipase nucleotide sequence.
- 15

Construction of expression plasmid pMT 2188.

- Aspergillus oryzae* expression plasmid pCaHj 483 (described in WO 98/00529) consists of an expression cassette based on the *Aspergillus niger* neutral amylase II promoter fused to the *Aspergillus nidulans* triose phosphate isomerase non translated leader sequence (Pna2/tpi) and the *Aspergillus niger* amyloglycosidase terminator (Tamg). Also present on the plasmid is the *Aspergillus* selective marker *amdS* from *Aspergillus nidulans* enabling growth on acetamide as
- 20

sole nitrogen source. These elements were cloned into the *E. coli* vector pUC19. The ampicillin resistance marker enabling selection in *E. coli* of this plasmid was replaced with the URA3 marker of *Saccharomyces cerevisiae* that can complement a *pyrF* mutation in *E. coli* in the following way:

- 5 The pUC19 origin of replication was PCR amplified from pCaHj483 with the primers 142779 and 142780 (SEQ ID NO: 40 and 41). The primer 142780 introduces a Bbu I site in the PCR fragment.

 The Expand PCR system (Roche Molecular Biochemicals, Basel, Switzerland) was used for the amplification following the manufacturers instructions
10 for this and the subsequent PCR amplifications.

 The URA3 gene was amplified from the general *S. cerevisiae* cloning vector pYES2 (Invitrogen corporation, Carlsbad, Ca, USA) using the primers 140288 and 142778 (SEQ ID NO: 36 and 39). The primer 140288 introduces an EcoR I site in the PCR fragment.

- 15 The two PCR fragments were fused by mixing them and amplifying using the primers 142780 and 140288 using the splicing by overlap method (Horton et al (1989) Gene, 77, 61-68).

 The resulting fragment was digested with EcoR I and Bbu I and ligated to the largest fragment of pCaHj 483 digested with the same enzymes. The ligation mixture
20 was used to transform the *pyrF E. coli* strain DB6507 (ATCC 35673) made competent by the method of Mandel and Higa (Mandel, M. and A. Higa (1970) J. Mol. Biol. 45, 154). Transformants were selected on solid M9 medium (Sambrook et. al (1989) Molecular cloning, a laboratory manual, 2. edition, Cold Spring Harbor Laboratory Press) supplemented with 1 g/l casaminoacids, 500 µg/l thiamine and 10
25 mg/l kanamycin.

 A plasmid from such a transformant was termed pCaHj 527.

 The Pna2/tpi promoter present on pCaHj527 was subjected to site directed mutagenesis by a simple PCR approach.

- Nucleotide 134 – 144 was altered from gtactaaaacc (SEQ ID NO: 57) to
30 ccgttaaattt (SEQ ID NO: 58) using the mutagenic primer 141223 (SEQ ID NO: 38).

 Nucleotide 423 – 436 was altered from atgcaatttaaact (SEQ ID NO: 59) to cggcaatttaacgg (SEQ ID NO: 60) using the mutagenic primer 141222: (SEQ ID NO: 37). The resulting plasmid was termed pMT 2188.

Expression of lipase gene in *Aspergillus oryzae*.

- 35 The plasmid pT152-1 was transformed to JM110 and non-methylated pT152-1 was extracted. The lipase gene was digested from non-methylated pT152-1 with Bcl I and Xho I and ligated into the BamH I and XhoI sites in the *Aspergillus*

expression cassette pMT2188 which has *Aspergillus niger* neutral amylase promoter, *Aspergillus nidulans* TPI leader sequences, *Aspergillus niger* glucoamylase terminator and *Aspergillus nidulans* amdS gene as a marker and *Saccharomyces cerevisiae* URA3 gene as a marker for a plasmid construction. The ligation mixture
5 was transformed E.coli 6507 by electroporation and the resultant plasmid was pNL152-3 (24).

pNL152-3 (24) was transformed into *Aspergillus oryzae* BECh-2. The selected transformants were inoculated in 100 ml of MS-9 media and cultivated at 30°C for 1 day. 3 ml of grown cell in MS-9 medium was inoculated to 100 ml of MDU-
10 2BP medium and cultivated at 32°C for 3 days. The supernatant was obtained by centrifugation.

The lipase productivity of selected transformants was determined as LU activity.

Example 15: Cloning and expression of lipase gene from *Fusarium culmorum*

15 Transformation in *Aspergillus* strain

Aspergillus oryzae strain BECh-2 was inoculated to 100 ml of YPG medium and incubated for 16 hrs at 32°C at 120 rpm. Pellets were collected and washed with 0.6 M KCl, and resuspended in 20 ml 0.6 M KCl containing a commercial β -glucanase product (Glucanex, product of Novo Nordisk A/S) at the concentration of 30 μ l/ml.
20 Cultures were incubated at 32°C at 60 rpm until protoplasts formed, then washed with STC buffer twice. The protoplasts were counted with a hematology counter and resuspended in an 8:2:0.1 solution of STC:STPC:DMSO to a final concentration of 2.5×10^7 protoplasts/ml. About 3 μ g of DNA was added to 100 μ l of protoplasts solution, mixed gently and incubated on ice for 30 min. One ml of STPC was added
25 and incubated 30 min at 37°C. After the addition of 10 ml of 50°C Cope top agarose, the reaction was poured onto Cope agar plate. Transformation plates were incubated at 32°C for 5 days.

PCR screening of lipase

A strain of *Fusarium culmorum* was used as a genomic DNA supplier.
30 PCR reactions on *Fusarium culmorum* genomic DNA was done with two following primer set: lip2 / lip21 (SEQ ID NO: 19/25) designed based upon the alignment of 3 lipases from *Fusarium*.

Reaction components (6 ng μ l of genomic DNA, 250 mM dNTP each, primer 250 nM each, 0.1 U μ l in Taq polymerase in 1X buffer (Roche Diagnostics, Japan))
35 were mixed and submitted for PCR under the following conditions.

Step	Temperature	Time
1	94°C	1 min
2	50°C	1 min
3	72°C	2 min
4	72°C	10 min
5	4°C	forever

Steps 1 to 3 were repeated 30 times.

0.7 kbp of fragment was amplified. It was gel-purified with GFX™ PCR DNA and Gel Band Purification kit (Amersham Pharmacia Biotech) and ligated into a pT7Blue vector with ligation high (TOYOBO, Japan). The ligation mixtures were transformed into *E. coli* JM109. The resultant plasmids, pT12-0221 was sequenced and compared to the *Fusarium oxysporum* lipase, showing that a clone encodes the internal part of the lipase.

Cloning of lipase gene

10 In order to clone the missing part of the lipase gene, LA PCR™ in vitro Cloning Kit (TaKaRa) was used for genome walking. 0.5 kbp of DNA fragment corresponding to N-terminal region was obtained from BamH I digested genome ligated to Sau3A I cassette of the kit with 12N1 primer (SEQ ID NO: 42). 1.8 kb of DNA fragment corresponding to C-terminal region was obtained from Bgl II digested
15 genome ligated to Sau3A I cassette of the kit with 12C2 primer (SEQ ID NO: 43).

Obtained fragments were purified by GFX™ PCR DNA and Gel Band Purification kit (amersham pharmacia biotech) and sequenced with each primer which amplified the fragment. Their sequence were compared to the *Fusarium oxysporum* lipase, showing that the amplified DNA covered N-terminal and C-
20 terminasl part of the lipase.

The fidelity of taq polymerase is not so good so in order to get the right sequence whole gene was amplified the primers 12-N (Bcl) and 12-C(Sal) (SEQ ID NO: 44 and 45).

Reaction components (6 ng / μ l of genomic DNA, 250 mM dNTP each, primer
25 250 nM each, 0.05 U/ μ l of Expand high fidelity polymerase in 1X buffer (Roche Diagnostics, Japan) were mixed and submitted for PCR under the following conditions.

Step	Temperature	Time
1	94°C	2 min
2	94°C	10sec
3	55°C	30sec
4	68°C	45sec
step 2-4 repeat 10 times		
5	94°C	10sec
6	55°C	30sec
7	68°C	45sec +20sec/cycle
step 5-7, repeat 20 times		
8	68°C	7min
7	4°C	forever

An amplified DNA fragment was gel-purified with GFX™ PCR DNA and Gel Band Purification kit (amersham pharmacia biotech) and ligated into a pT7Blue vector with ligation high (TOYOBO, Japan). The ligation mixtures were transformed into *E. coli* JM109. The resultant plasmids, pT12-1, pT12-2, pT12-3, and pT12-4, were sequenced and all of them are identical. The sequence is defined as *Fusarium culmorum* lipase DNA sequence.

Expression of lipase gene in *Aspergillus oryzae*.

The plasmid pT12-6 was transformed to JM110 and non-methylated pT12-6 was extracted. The lipase gene was digested from non-methylated pT12-6 with Bcl I and Sal I into T-vector and ligated into the BamH I and XhoI sites in the *Aspergillus* expression cassette pMT2188 which has *Aspergillus niger* neutral amylase promoter, *Aspergillus nidulans* TPI leader sequences, *Aspergillus niger* glucoamylase terminator and *Aspergillus nidulans* amdS gene as a marker and *Saccharomyces cerevisiae* URA3 gene as a marker for a plasmid construction. The ligation mixture was transformed *E. coli* 6507 by electroporation and the resultant plasmid was pNL12-***.

pNL12-*** was transformed into *Aspergillus oryzae* BECh-2. The selected transformants were inoculated in 100 ml of MS-9 media and cultivated at 30°C for 1 day. 3 ml of grown cell in MS-9 medium was inoculated to 100 ml of MDU-2BP medium and cultivated at 32°C for 3 days. The supernatant was obtained by centrifugation.

The lipase productivity of selected transformants was determined as LU activity.

Example 16: Cloning of a phospholipase gene from the *Fusarium solani* strain MUCL 38667.

5 A genomic DNA preparation of the strain MUCL 38667 was made as described in WO 00/24883.

A PCR reaction (96°C 5 min, 30* (94°C 30 sec., 55°C 30 sec, 72°C 1 min), 72°C 5 min) was run using PWO polymerase in 2.5 mM MgSO₄ as recommended by the manufacturer (Boehringer Mannheim) with the MUCL 38667 genomic DNA as
10 template, with oligo 161000J1 and 161000J2 (SEQ ID NO: 46 and 47). These oligo'es were designed based conserved sequences in homologous phospholipases.

A fragment of 180 bp was isolated from a 2 % gel. Because the amounts of DNA was very small, a new identical pcr was run, this time using the 180 bp
15 fragment as template rather than MUCL 38667 genomic DNA.

This fragment was cloned into pCR4 using TOPO-cloning as recommended by the manufacture (Invitrogen) and transformed into the *E. coli* strain TOPO10.

DNA preparations where made using the Qiagen minispinprep kit and the clones where sequenced using M13 rev and M13 fwp primer supplied with the
20 TOPO-cloning kit (Invitrogen). Sequence alignment was made to all available DNA sequence using SRS. The 180 bp fragment was identified as originating from a phospholipase gene.

Based on the 180 bp fragment DNA sequence, four primers were designed: 071200J1; 071200j2, 221200J1, 221200J2 (SEQ ID NO: 48-51).

25 The MUCL38667 genomic DNA (app. 1 µg) was cut with AgeI in a volume of 10 µl and ligated in a volume of 500 µl. The DNA was precipitated in ethanol and redissolved in water. 2 µl of the religated mix was used as template, and oligo 071200J1 and 071200J2 in a PCR reaction using GeneAMP XL PCR kit as recommended by manufacture (Boehringer Mannheim) in a total of 20 µl.

30 1 µl of this PCR reaction fragments was used as template in a second PCR reaction using nested oligoes 221200J1 and 221200J2 (SEQ ID NO: 50 and 51), which was identical to the above mentioned.

The generated PCR fragment of app. 1500 bp was cloned into pCR4 using TOPO-cloning as recommended by the manufacture (Invitrogen) and transformed
35 into the *E. coli* strain TOPO10.

DNA preparations were made using the Qiagen minispinprep kit and the clones were sequenced using M13 rev and M13 fwd primer supplied with the TOPO-cloning kit (Invitrogen). Sequence alignment was made to all available DNA sequence using SRS. The 1500 bp fragment was identified as originating from the 3' end of a phospholipase gene.

Based on the 1500 bp fragment DNA sequence, one primer was designed: 170101J11 (SEQ ID NO: 52).

The MUCL 38667 genomic DNA (app. 1 µg) was cut with HindIII in a volume of 10 µl and ligated in a volume of 500 µl. The DNA was precipitated in ethanol and redissolved in water. 2 µl of the religated mix was used as template, and oligo 221200J1 and 170101J11 in a PCR reaction using GeneAMP XL PCR kit as recommended by manufacture (Boehringer Mannheim) in a total of 20 µl.

The generated PCR fragment of app. 350 bp was cloned into pCR4 using TOPO-cloning as recommended by the manufacture (Invitrogen) and transformed into the *E. coli* strain TOPO10.

DNA preparations were made using the Qiagen minispinprep kit and the clones were sequenced using T3 and T7 primer supplied with the TOPO-cloning kit (Invitrogen). Sequence alignment was made to all available DNA sequence using SRS. The 350 bp fragment was identified as originating from the 5' end of a phospholipase gene.

Based on the 350 bp and the 1500 bp DNA sequence, two primers were designed (290101J2 and 020301J1, SEQ ID NO: 53 and 54), thus covering the whole gene.

A PCR reaction (96°C 5 min, 30* (94°C 30 sec., 55°C 30 sec, 72°C 2 min), 72°C 5 min) was run using PWO polymerase in 2.5 mM MgSO₄ as recommended by manufacture (Boehringer Mannheim) with the MUCL38667 genomic DNA as template, with oligo 290101J2 and 020301J1 (SEQ ID NO: 53 and 54).

The generated PCR fragment of app. 1100 bp were cloned into pCR4 using TOPO-cloning as recommended by the manufacture (Invitrogen) and transformed into the *E. coli* strain TOPO10.

DNA preparations were made using the Qiagen minispinprep kit and the clones were sequenced using T3 and T7 primer supplied with the TOPO-cloning kit (Invitrogen). Sequence alignment was made to all available DNA sequence using SRS, as well as earlier sequence of the same.

Example 17: Cloning and expression of lipase gene from *Fusarium solani*

The cloned phospholipase gene in the TOPO vector, as well as pJVI9 (WO 97/47746) was cut with the restriction enzymes BamHI and XhoI. The pJVI9 vector and the phospholipase gene were purified from a 1% agarose gel, and ligated o/n.

5 The ligation was transformed into the *E.coli* strain DH10b, and transformants were isolated.

DNA preparations were made using the Qiagen minispinprep kit and the clones were verified by sequencing using 19670 and 19671 primer (SEQ ID NO: 55 and 56).

10 The resulting plasmid was transformed into the *Aspergillus oryzae* strain Jal125 (WO 97/35956) using the following method:

Transformation of *Aspergillus oryzae* (general procedure)

100 ml of YPD (Sherman et al., (1981), Methods in Yeast Genetics, Cold Spring Harbor Laboratory) are inoculated with spores of *A. oryzae* and incubated
15 with shaking for about 24 hours. The mycelium is harvested by filtration through miracloth and washed with 200 ml of 0.6 M MgSO₄. The mycelium is suspended in 15 ml of 1.2 M MgSO₄, 10 mM NaH₂PO₄, pH 5.8. The suspension is cooled on ice and 1 ml of buffer containing 120 mg of Novozym[®] 234 is added. After 5 min., 1 ml of 12 mg/ml BSA (Sigma type H25) is added and incubation with gentle agitation
20 continued for 1.5-2.5 hours at 37°C until a large number of protoplasts is visible in a sample inspected under the microscope.

The suspension is filtered through miracloth, the filtrate transferred to a sterile tube and overlaid with 5 ml of 0.6 M sorbitol, 100 mM Tris-HCl, pH 7.0. Centrifugation is performed for 15 min. at 1000 g and the protoplasts are collected
25 from the top of the MgSO₄ cushion. 2 volumes of STC (1.2 M sorbitol, 10 mM Tris-HCl, pH 7.5, 10 mM CaCl₂) are added to the protoplast suspension and the mixture is centrifuged for 5 min. at 1000 g. The protoplast pellet is resuspended in 3 ml of STC and repelleted. This is repeated. Finally, the protoplasts are resuspended in 0.2-1 ml of STC.

30 100 µl of protoplast suspension are mixed with 5-25 µg of p3SR2 (an *A. nidulans* amdS gene carrying plasmid described in Hynes et al., Mol. and Cel. Biol., Vol. 3, No. 8, 1430-1439, Aug. 1983) and 5 µg of the pJVI9-phospholipase plasmid in 10 µl of STC. The mixture is left at room temperature for 25 min. 0.2 ml of 60% PEG 4000 (BDH 29576), 10 mM CaCl₂ and 10 mM Tris-HCl, pH 7.5 is added and
35 carefully mixed (twice) and finally 0.85 ml of the same solution are added and carefully mixed. The mixture is left at room temperature for 25 min., spun at 2,500 g for 15 min. and the pellet is resuspended in 2 ml of 1.2M sorbitol. After one more

sedimentation the protoplasts are spread on minimal plates (Cove, (1966), Biochem. Biophys. Acta 113, 51-56) containing 1.0 M sucrose, pH 7.0, 10 mM acetamide as nitrogen source and 20 mM CsCl to inhibit background growth. After incubation for 4-7 days at 37°C spores are picked, suspended in sterile water and spread for single colonies.

12 independent transformants from the pJV19-phospholipase transformations were isolated on minimal plates (Cove, (1966), Biochem. Biophys. Acta 113, 51-56) containing 1.0 M sucrose, pH 7.0, 10 mM acetamide as nitrogen source and 20 mM CsCl to inhibit background growth, and at the same time inoculated into a 96-well microtiter dish containing 200 µl minimal media of 1% Vogel, 2% maltose (e.g., Methods in Enzymology, Vol. 17 p. 84) in each well.

After three days of incubation at 34°C, media from the cultures in the microtiter dish were assayed for lipase activity. A 10 µl aliquot of media from each well was added to a microtiter well containing 200 µl of a lipase substrate of 0.018% p-nitrophenylbutyrate, 0.1% Triton X-100, 10 mM CaCl₂, 50 mM Tris pH 7.5. Activity was assayed spectrophotometrically at 15-second intervals over a five minute period, using a kinetic microplate reader (Molecular Device Corp., Sunnyvale CA), using a standard enzymology protocol (e.g., *Enzyme Kinetics*, Paul C. Engel, ed., 1981, Chapman and Hall Ltd.) Briefly, product formation is measured during the initial rate of substrate turnover and is defined as the slope of the curve calculated from the absorbance at 405 nm every 15 seconds for 5 minutes.

This procedure was repeated and spores of the best producing transformants after the second re-isolation were stored as a defined transformant.

A. oryzae JaL 125 (WO 97/35956) is derived from *Aspergillus oryzae* IFO 4177 available from Institute for Fermentation, Osaka; 17-25 Juso Hammachi 2-Chome Yodogawa-ku, Osaka, Japan, having the alkaline protease gene named "alp" (described by Murakami K et al., (1991), Agric. Biol. Chem. 55, p. 2807-2811) deleted by a one step gene replacement method (described by G. May in "Applied Molecular Genetics of Filamentous Fungi" (1992), p. 1-25. Eds. J. R. Kinghorn and G. Turner; Blackie Academic and Professional), using the *A. oryzae* pyrG gene as marker.

Example 18: Preparation and testing of variant

An R275A variant of *Acremonium berkeleyanum* lipase (SEQ ID NO: 6) was constructed and expressed in *A. oryzae*. Preliminary tests using crude culture broth confirmed the variant enzyme to be more heat stable than the wildtype. SDS-PAGE showed that around 60% of the molecules still were processed, compared to 100%

for the wildtype. The temperature-stability curve of the crude variant broth was found to be biphasic, and the unprocessed form seemed to be around 10°C more stable than the processed one.

Two bands were seen in SDS-PAGE: unprocessed lipase at around 31 kDa and C-terminal processed at ca. 28 kDa. It was found that unprocessed lipase is stable for 30 minutes up until ca. 60°C, whereas the wildtype, processed lipase is degraded quickly at temperature over 50°C.

The above findings indicate that the R275A variant of *Acremonium berkeleyanum* lipase is to a certain degree protected against KEX2 type protease attack in *Aspergillus oryzae*.

Example 19: Determination of phospholipase and galactolipase activities

The lipolytic enzymes from *F. venenatum*, *F. sulphureum*, *A. berkeleyanum*, *F. culmorum* and *F. solani* (SEQ ID NO: 2, 4, 6, 8 and 10) were tested for phospholipase and galactolipase activities by plate assay methods, as described in WO 0032758. The lipolytic enzyme from *F. oxysporum* was included for comparison.

The results showed that all 5 enzymes have phospholipase activity. The enzymes from *F. venenatum*, *F. sulphureum* and *F. culmorum* and *F. oxysporum* have activities at a similar level, and the other two have slightly lower activities.

All 5 enzymes were found to have galactolipase activity (digalactosyl diglyceride hydrolyzing activity). The enzymes from *F. venenatum*, *F. sulphureum*, *F. culmorum* and *F. oxysporum* have significantly higher activity than those from *A. berkeleyanum* and *F. solani*.

Example 20: Preparation of straight-dough bread

Doughs were made using spiral mixers from 2 kg of Meneba flour (batch 941-2) according to the straight dough method (AACC Method 10-10B in Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

Lipase activity was determined using pH-stat titration and tributyrin as substrate at 30°C and pH 7. Enzymes were dosed according to the protocol below:

30

35

Dough	1	2	3	4	5	6	7	8
Enzyme								
<i>F. venenatum</i> Lipase, LU/kg flour			10	20	40	80	100	150
<i>F. oxysporum</i> Lipase, LU/kg flour		1000						
Fungamyl, FAU/kg flour	10	10	10	10	10	10	10	10
PentMono, FXU/kg flour	100	100	100	100	100	100	100	100
DATEM, %	0.4							

Example 21: Dough and bread evaluation and dough stability measurements

Each dough described in the previous Example was split into 15 rolls for 45 minute fermentation; 15 rolls for 70 minute fermentation. Two pan breads were evaluated for of the breads crumb structure; and 2 pan breads were evaluated for 5 texture analysis.

Firmness, elasticity of the doughs, and crumb structure, texture of the breads, shape factor and volume of the rolls were measured using the methods described earlier. Bread was measured 2 and 24 hours after baking.

The results of Dough 5 showed that the effect of the *Fusarium venenatum* lipase dosed at 40 LU/kg flour was similar to that of the *Fusarium oxysporum* lipase 10 dosed at 1000 LU/kg flour. Increased dosages of the *Fusarium venenatum* lipase resulted in softer, more sticky dough which had a lower extensibility and higher elasticity. At a dosage of 150 LU/kg flour, the *Fusarium venenatum* lipase yielded a dough which broke apart easily when stretched, similar to undermixed doughs.

15 The effect of the *Fusarium venenatum* lipase on dough parameters was dosage dependent. Generally bread made from dough treated with the *Fusarium venenatum* lipase yielded a more uniform crumb, a coarser grain with thicker cell walls and less elongated shape, and a less white crumb colour compared to *Fusarium oxysporum* lipase.

20 The effect of the lipases on stability is shown in Tables 1 and 2 at 45 minutes and 70 minutes, respectively. After normal fermentation time, addition of the *Fusarium venenatum* lipase yielded a very good stability factor when dosed at and above 80 LU/kg flour. At dosages of 100 and 150 LU of the *Fusarium venenatum*

lipase per kg flour, the volume of the bread matched that of bread treated with DATEM (di-acetylated-tartaric acid-esters of mono-and diglycerides of fatty acids), but was slightly smaller than that of the bread treated with the *Fusarium oxysporum* lipase. The best shape-factors were obtained with the high dosages of the *Fusarium*
 5 *venenatum* lipase.

Table 1

Lipolytic enzyme added	Bread	
	Sp. Vol. (ml/g)	Shape factor
	45 min	45 min
Datem	6.89	0.672
F. oxysporum (200LU)	7.18	0.696
F. venenatum (10LU)	6.34	0.663
F. venenatum (20LU)	6.64	0.681
F. venenatum (40LU)	6.45	0.653
F. venenatum (80LU)	6.51	0.687
F. venenatum (100LU)	6.75	0.716
F. venenatum (150LU)	6.85	0.700

Table 2

Lipolytic enzyme added	Bread	
	Sp. Vol. (ml/g)	Shape factor
	70 min	70 min
Datem	8.11	0.652
F. oxysporum (200LU)	7.96	0.682
F. venenatum (10LU)	6.76	0.648
F. venenatum (20LU)	6.79	0.655
F. venenatum (40LU)	6.94	0.651
F. venenatum (80LU)	7.54	0.672
F. venenatum (100LU)	7.69	0.678
F. venenatum (150LU)	7.78	0.680

After extended fermentation time, 80 – 150 LU/kg flour of the *Fusarium venenatum* lipase yielded a better shape-factor than DATEM, and 150 LU/kg flour of the *Fusarium venenatum* lipase performed as well as the *Fusarium oxysporum* lipase. A larger volume was achieved with increased dosage of the *Fusarium*
5 *venenatum* lipase, where at 150 LU of the lipase per kg of flour, the volume increase nearly matched the volume increase with the *Fusarium oxysporum* lipase and DATEM.

When dosed at 100 and 150 LU/kg flour, the *Fusarium venenatum* lipase yielded a better shape-factor, but lower volume after normal fermentation time than
10 the *Fusarium oxysporum* lipase. After an extended fermentation time, 100 and 150 LU/kg flour of *Fusarium venenatum* lipase performed as well as the *Fusarium oxysporum* lipase.

Firmness and elasticity were followed during the first 24 hours after baking. The *Fusarium venenatum* lipase performed similarly to the *Fusarium oxysporum*
15 lipase on initial softness and elasticity.

The overall optimal dosage of the *Fusarium venenatum* lipase was between 80–150 LU/kg flour, which corresponded to approximately 0.4–0.7 mg enzyme/kg flour. The optimal performance of *Fusarium venenatum* lipase appeared to be very close to that of the *Fusarium oxysporum* lipase at the optimal dosage of 1000 LU per
20 kg flour corresponding to 0.33 mg enzyme/kg flour.

Original (for SUBMISSION) - printed on 25.06.2001 03:39:58 PM

0-1	Form - PCT/RO/134 (EASY) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared using	PCT-EASY Version 2.91 (updated 01.01.2001)
0-2	International Application No.	
0-3	Applicant's or agent's file reference	10055-WO
1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	3-4
1-2	line	23-5
1-3	Identification of Deposit	
1-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
1-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
1-3-3	Date of deposit	15 June 2000 (15.06.2000)
1-3-4	Accession Number	DSMZ 13539
1-4	Additional Indications	NONE
1-5	Designated States for Which Indications are Made	all designated States
1-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
2	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
2-1	page	3-4
2-2	line	23-5
2-3	Identification of Deposit	
2-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
2-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
2-3-3	Date of deposit	15 June 2000 (15.06.2000)
2-3-4	Accession Number	DSMZ 13538
2-4	Additional Indications	NONE
2-5	Designated States for Which Indications are Made	all designated States
2-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE

Original (for SUBMISSION) - printed on 25.06.2001 03:39:58 PM

3	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
3-1	page	3-4
3-2	line	23-5
3-3	Identification of Deposit	
3-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
3-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
3-3-3	Date of deposit	15 June 2000 (15.06.2000)
3-3-4	Accession Number	DSMZ 13537
3-4	Additional Indications	NONE
3-5	Designated States for Which Indications are Made	all designated States
3-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
4	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
4-1	page	3-4
4-2	line	23-5
4-3	Identification of Deposit	
4-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
4-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
4-3-3	Date of deposit	21 June 2001 (21.06.2001)
4-3-4	Accession Number	DSMZ 14361
4-4	Additional Indications	NONE
4-5	Designated States for Which Indications are Made	all designated States
4-6	Separate Furnishing of Indications These indications will be submitted to the International Bureau later	NONE
5	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
5-1	page	3-4
5-2	line	23-3

Original (for SUBMISSION) - printed on 25.06.2001 03:39:58 PM

5-3	Identification of Deposit	
5-3-1	Name of depositary institution	Agricultural Research Service Culture Collection
5-3-2	Address of depositary institution	1815 North University Street, Peoria, Illinois 61604 ,United States of America
5-3-3	Date of deposit	22 August 2000 (22.08.2000)
5-3-4	Accession Number	NRRL B-30333
5-4	Additional Indications	Deposit resubmitted on 30 March 2001
5-5	Designated States for Which Indications are Made	all designated States
5-6	Separate Furnishing of Indications	NONE
	These indications will be submitted to the International Bureau later	

FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	yes
0-4-1	Authorized officer	<i>Mike C. DeBaker</i>

FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	

CLAIMS

1. A polynucleotide which comprises a nucleotide sequence encoding a lipolytic enzyme, which sequence:
 - a) is a DNA sequence cloned into a plasmid present in *Escherichia coli* deposit number *Escherichia coli* NRRL B-30333, DSM 13537, DSM 13538, DSM 13539 or DSM 14361,
 - b) is the mature polypeptide-encoding DNA sequence shown in SEQ ID NO: 1, 3, 5, 7 or 9 or is a DNA sequence that can be derived therefrom by substitution, deletion, and/or insertion of one or more nucleotides, or
 - c) has at least 80 % identity with the mature polypeptide-encoding sequence of SEQ ID NO: 1, 3 or 7, or at least 70 % identity with the mature polypeptide-encoding sequence of SEQ ID NO: 5 or 9; or
 - d) a complementary strand of the sequence defined in a), b) or c).
2. The polynucleotide of claim 1 which is native to a strain of *Fusarium* or *Acremonium*, particularly *F. venenatum*, *F. sulphureum*, *A. berkeleyanum*, *F. culmorum* or *F. solani*.
3. A polypeptide having lipolytic enzyme activity which is:
 - a) a polypeptide having an amino acid sequence which has at least 92 % identity with the mature polypeptide of SEQ ID NO: 2, at least 85 % identity with the mature polypeptide of SEQ ID NO: 4 or at least 70 % identity with the mature polypeptide of SEQ ID NO: 6;
 - b) a polypeptide which is encoded by a nucleic acid sequence which hybridizes under high stringency conditions with a complementary strand of the nucleic acid sequence encoding the mature polypeptide of SEQ ID NO: 1, 3 or 5 or a subsequence thereof having at least 100 nucleotides;
 - c) a polypeptide having an amino acid sequence which can be obtained from the mature polypeptide of SEQ ID NO: 2, 4 or 6 by substitution, deletion, and/or insertion of one or more amino acids; or
 - d) a polypeptide encoded by the lipolytic enzyme encoding part of the DNA sequence cloned into a plasmid present in *Escherichia coli* deposit number NRRL B-30333, DSM 13538 or DSM 13539, or DSM 14361

4. The lipolytic enzyme of claim 3 which is native to a strain of *Fusarium* or *Acremonium*, particularly *F. venenatum*, *F. sulphureum* or *A. berkeleyanum*.
5. The lipolytic enzyme of claim 3 or 4 which has an amino acid sequence comprising the mature polypeptide of SEQ ID NO: 2, 4 or 6.
- 5 6. A polynucleotide comprising a nucleic acid sequence which encodes the lipolytic enzyme of any of claims 3-5.
7. A nucleic acid construct comprising the polynucleotide of claim 1, 2 or 6 operably linked to one or more control sequences capable of directing the expression of the lipolytic enzyme in a suitable expression host.
- 10 8. A recombinant expression vector comprising the nucleic acid construct of claim 7, a promoter, and transcriptional and translational stop signals.
9. A recombinant host cell comprising the nucleic acid construct of claim 7.
10. A method for producing a lipolytic enzyme comprising cultivating the host cell of claim 10 under conditions conducive to production of the lipolytic enzyme, and
15 recovering the lipolytic enzyme.
11. A method for producing a mutant nucleic acid sequence, comprising
 - a) introducing at least one mutation into the mature polypeptide coding sequence of SEQ ID NO: 1, 3, 5, 7 or 9, and
 - b) recovering the mutant nucleic acid sequence.
- 20 12. A method for producing a polypeptide comprising
 - a) introducing at least one mutation into a polynucleotide having the mature polypeptide coding sequence of SEQ ID NO: 1, 3, 5, 7 or 9,
 - b) transforming a host cell with the mutated polynucleotide,
 - c) cultivating the transformed host cell under conditions conducive for
25 production of polypeptides,
 - d) screening the polypeptides to select a polypeptide having lipolytic enzyme activity, and
 - e) producing the selected polypeptide.

13. A method of producing a lipolytic enzyme which comprises:
- a) shuffling a first polynucleotide comprising the mature polypeptide coding sequence of SEQ ID NO: 1, 3, 5, 7 or 9 and a second polynucleotide encoding a lipolytic enzyme and having at least 50 % identity to the first polynucleotide,
 - b) expressing the shuffled polynucleotides to form recombinant polypeptides,
 - c) screening the polypeptides to select a polypeptide having lipolytic enzyme activity, and
 - d) producing the selected polypeptide.
- 10 14. A method for preparing a dough or a baked product made from the dough, comprising adding the lipolytic enzyme of any of claims 3-5 to the dough.
15. A dough composition comprising the lipolytic enzyme of any of claims 3-5.
16. A detergent composition comprising a surfactant and the lipolytic enzyme of claim any of claims 3-5.

1/2




Fig. 1

2/2




Fig. 2

10055-WO.ST25
SEQUENCE LISTING

<110> Novozymes

<120> Lipolytic enzymes and genes encoding them

<130> 10055-WO

<160> 60

<170> PatentIn version 3.0

<210> 1

<211> 2940

<212> DNA

<213> *Fusarium venenatum*

<220>

<221> CDS

<222> (1376)..(1465)

<220>

<221> CDS

<222> (1515)..(1681)

<220>

<221> CDS

<222> (1740)..(2529)

<220>

<221> mat_peptide

<222> (1515)..()

<400> 1

aattcatgtg aatctactat gtaacagtat gttgtattgc attacccatc aacgttgaat

60

10055-WO.ST25

cgttgcgacg taacggcccc gttcaagcga gatgtagata tgttggtagt taattgatgg	120
gttaggtatt cttttcatca actcgggtatt ctcatccccc agatatcggc acttgtcttt	180
actccagatt tcatatcgca tcgagttata tacagtccca attgagtcga ctaccccgtc	240
caaaacaggt tttctcacia accaaccgca gcctaacaaa aagtcctttg tctttctgca	300
ataaatgctg acaccccctg gcttttttagg actgacggct cacgatgcag ccgttgcgat	360
aattaattga caattacccg cacattgatg catacttggc ggtcagggtca ggtcaggctg	420
aagcatacct attgggtcat ttatttgccg atcgtggtga aaagaatgca agtgataact	480
agttacgagt cgctttatga aagatgggtg gtcgaaactg tcaatatggc atgggaggca	540
aatcgtttgg tctcaactct atagcatgta ctataattgg tcttttcatc acagtcacgc	600
caaagtgccg gtctcagact atggaccaac cactttcctc cttcacgtct aaattgactt	660
gatcaccaga ctgcaatatt ttttcttttc ttctataccc ctaggatcat acaatacgaa	720
ccccaaactca actcgagaga gagagtcccc ttcccaacat ttgacagcc cttgctcttc	780
tcctccagg atgtaacaga agctgaaagg gtacccctgt agcccacctt taccacccat	840
cttttccatc tgtatcgggt catcccatca caaccctcac gtggtccgag atcgtcgta	900
cccgtattgg aagctcactc cgggcccaac gagagattgg accaaggaaa aataactttg	960
agacctcttc aagcagtcgg tcattcgta ctgggatgtg tagtcgataa tgcggggtga	1020
caggccctca atccagcacc caccatcatg ggcactgact gtactaccgg agcccatcat	1080
ttcgtttttg ggtcctggcg tctacttgac cgactgagtt tgccaagatg gatggcatga	1140
gagacagtgg ttaggctggg cgggtattgt gatgagagaa agcgagagac tagttagaag	1200
caaagaaaaa agatatataa gctgtcacat ccctcatgaa catgctgttc ttgtaagtcg	1260
ggatatcagg gccagcttca gtattcagta tcctttctga gggagttgca cttgtcaca	1320
gcttgtctgt ctatcactta tacttaccct tggaccacgt tctttgtctg tcaag atg	1378
	Met -30
cat ctc cta tca ctc ctc tca att gcc acc ctt gcg gta gcc agc cct	1426
His Leu Leu Ser Leu Leu Ser Ile Ala Thr Leu Ala Val Ala Ser Pro	-25 -20 -15
ctg agc gtt gaa gat tac gcc aag gct ctc gat gaa aga ggtaaaacga	1475
Leu Ser Val Glu Asp Tyr Ala Lys Ala Leu Asp Glu Arg	-10 -5 -1
ttctctgttc ccataacaat tccaatactc acagaccta gct gtt tct gtc tct	1529
	Ala Val Ser Val Ser 1 5
acc aac gac ttt ggc aac ttc aag ttc tac atc cag cac ggt gcc gca	1577
Thr Asn Asp Phe Gly Asn Phe Lys Phe Tyr Ile Gln His Gly Ala Ala	10 15 20
gca tac tgt aac tct gaa gcc gca gcc ggt gca aag gtc acc tgc gga	1625
Ala Tyr Cys Asn Ser Glu Ala Ala Ala Gly Ala Lys Val Thr Cys Gly	25 30 35
gga aac ggt tgc cca acg gtc cag tcc aat ggt gcc acc atc gtg gca	1673
Gly Asn Gly Cys Pro Thr Val Gln Ser Asn Gly Ala Thr Ile Val Ala	40 45 50

10055-WO.ST25

tct ttc ct gtaagtctaa catatcaciaa acacatcatc aactccaaac	1721
Ser Phe Leu 55	
ttacaaatct ctttatag t ggc tca aag act ggc atc ggt ggc tac gtc gcg	1773
Gly Ser Lys Thr Gly Ile Gly Gly Tyr Val Ala 60 65	
acc gac tct gca cgc aag gaa atc gtc ctc tcg gtt cgc ggt agc acc	1821
Thr Asp Ser Ala Arg Lys Glu Ile Val Leu Ser Val Arg Gly Ser Thr 70 75 80	
aac att cgc aac tgg ctt acc aac ctc gac ttc gac cag gat gac tgc	1869
Asn Ile Arg Asn Trp Leu Thr Asn Leu Asp Phe Asp Gln Asp Asp Cys 85 90 95	
agc ttg acc tcc ggc tgt gga gtg cac gga ggc ttc cag aga gcc tgg	1917
Ser Leu Thr Ser Gly Cys Gly Val His Gly Gly Phe Gln Arg Ala Trp 100 105 110 115	
aat gag atc tcg gcc gca gca acc gcc gct gtc gca aag gcc cgc aag	1965
Asn Glu Ile Ser Ala Ala Ala Thr Ala Val Ala Lys Ala Arg Lys 120 125 130	
gcg aac cct tcg ttc aag gtc gtt gcc aca ggt cac tcg ttg ggt ggt	2013
Ala Asn Pro Ser Phe Lys Val Val Ala Thr Gly His Ser Leu Gly Gly 135 140 145	
gct gta gct aca cta gca ggt gcg aac ctg cga gtt ggt ggt acg cca	2061
Ala Val Ala Thr Leu Ala Gly Ala Asn Leu Arg Val Gly Gly Thr Pro 150 155 160	
gtt gac atc tac acc tac ggc tca ccc cga gtt gga aac acg caa ctc	2109
Val Asp Ile Tyr Thr Tyr Gly Ser Pro Arg Val Gly Asn Thr Gln Leu 165 170 175	
gct gcc ttc atc tct aac cag gct ggt gga gag ttc cgc gtt acg aac	2157
Ala Ala Phe Ile Ser Asn Gln Ala Gly Gly Glu Phe Arg Val Thr Asn 180 185 190 195	
gcc aag gac ccc gtg cct cgt ctc ccc cct ctg gtc ttt gga tac cgg	2205
Ala Lys Asp Pro Val Pro Arg Leu Pro Pro Leu Val Phe Gly Tyr Arg 200 205 210	
cac aca tcc ccc gag tac tgg ttg tct ggt agc gga ggt aac aag gtt	2253
His Thr Ser Pro Glu Tyr Trp Leu Ser Gly Ser Gly Gly Asn Lys Val 215 220 225	
gac tac acc atc aat gat gtc aag gtg tgt gag ggt gct gcc aac ctt	2301
Asp Tyr Thr Ile Asn Asp Val Lys Val Cys Glu Gly Ala Ala Asn Leu 230 235 240	
cag tgc aac ggt gga aca ctc gga ttg gat atc gac gcc cat ctc cac	2349
Gln Cys Asn Gly Gly Thr Leu Gly Leu Asp Ile Asp Ala His Leu His 245 250 255	
tac ttc cag gag acc gat gct tgc tct ggt tcc ggt atc gcg tgg aga	2397
Tyr Phe Gln Glu Thr Asp Ala Cys Ser Gly Ser Gly Ile Ala Trp Arg 260 265 270 275	
aga tac agg agt gct aag cgt gag agc atc tcg gag agg gcc act atg	2445
Arg Tyr Arg Ser Ala Lys Arg Glu Ser Ile Ser Glu Arg Ala Thr Met 280 285 290	
aca gat gcc gag ctg gag aag aag ctt aac aac tat gtt gcg atg gat	2493
Thr Asp Ala Glu Leu Glu Lys Lys Leu Asn Asn Tyr Val Ala Met Asp 295 300 305	

10055-WO.ST25

aag gag tat gtc aag act cac gcc aac cgc tca tcg tagtatgaca 2539
 Lys Glu Tyr Val Lys Thr His Ala Asn Arg Ser Ser
 310 315

tttacgcagt aacgatataa ttaccataac aaaaactctg gataccattc tgggtgcaagc 2599
 atggcgaaga aaacatcatt atctatgtga atgtatcata accatcctta cgccatgccg 2659
 ttgatcttac tactgagaca aaatactcag tcatgtacaa caaactccaa agcaccgaat 2719
 gacttctggc tttttggcaa agcacgaaac caatcattca aaccctcca cgaccatgcc 2779
 ctgcgcatgg ggaacacca cgagaatgac accacgaggc acgcggacac tcttcacctt 2839
 catgcacca aagacattga cttcccgat attagggcat gctcggaaaa tggaaccag 2899
 aacaaaatcc gtcactgcct cacagaaact gatctccaat t 2940

<210> 2

<211> 349

<212> PRT

<213> Fusarium venenatum

<400> 2

Met His Leu Leu Ser Leu Leu Ser Ile Ala Thr Leu Ala Val Ala Ser
 -30 -25 -20 -15

Pro Leu Ser Val Glu Asp Tyr Ala Lys Ala Leu Asp Glu Arg Ala Val
 -10 -5 -1 1

Ser Val Ser Thr Asn Asp Phe Gly Asn Phe Lys Phe Tyr Ile Gln His
 5 10 15

Gly Ala Ala Ala Tyr Cys Asn Ser Glu Ala Ala Ala Gly Ala Lys Val
 20 25 30

Thr Cys Gly Gly Asn Gly Cys Pro Thr Val Gln Ser Asn Gly Ala Thr
 35 40 45 50

Ile Val Ala Ser Phe Leu Gly Ser Lys Thr Gly Ile Gly Gly Tyr Val
 55 60 65

Ala Thr Asp Ser Ala Arg Lys Glu Ile Val Leu Ser Val Arg Gly Ser
 70 75 80

Thr Asn Ile Arg Asn Trp Leu Thr Asn Leu Asp Phe Asp Gln Asp Asp
 85 90 95

Cys Ser Leu Thr Ser Gly Cys Gly Val His Gly Gly Phe Gln Arg Ala
 100 105 110

Trp Asn Glu Ile Ser Ala Ala Ala Thr Ala Ala Val Ala Lys Ala Arg
 115 120 125 130

10055-WO.ST25

Lys Ala Asn Pro Ser Phe Lys Val Val Ala Thr Gly His Ser Leu Gly
 135 140 145

Gly Ala Val Ala Thr Leu Ala Gly Ala Asn Leu Arg Val Gly Gly Thr
 150 155 160

Pro Val Asp Ile Tyr Thr Tyr Gly Ser Pro Arg Val Gly Asn Thr Gln
 165 170 175

Leu Ala Ala Phe Ile Ser Asn Gln Ala Gly Gly Glu Phe Arg Val Thr
 180 185 190

Asn Ala Lys Asp Pro Val Pro Arg Leu Pro Pro Leu Val Phe Gly Tyr
 195 200 205 210

Arg His Thr Ser Pro Glu Tyr Trp Leu Ser Gly Ser Gly Gly Asn Lys
 215 220 225

Val Asp Tyr Thr Ile Asn Asp Val Lys Val Cys Glu Gly Ala Ala Asn
 230 235 240

Leu Gln Cys Asn Gly Gly Thr Leu Gly Leu Asp Ile Asp Ala His Leu
 245 250 255

His Tyr Phe Gln Glu Thr Asp Ala Cys Ser Gly Ser Gly Ile Ala Trp
 260 265 270

Arg Arg Tyr Arg Ser Ala Lys Arg Glu Ser Ile Ser Glu Arg Ala Thr
 275 280 285 290

Met Thr Asp Ala Glu Leu Glu Lys Lys Leu Asn Asn Tyr Val Ala Met
 295 300 305

Asp Lys Glu Tyr Val Lys Thr His Ala Asn Arg Ser Ser
 310 315

<210> 3

<211> 1161

<212> DNA

<213> Fusarium sulphureum

<220>

<221> CDS

<222> (1)..(94)

<220>

10055-WO.ST25

<221> CDS

<222> (147)..(312)

<220>

<221> CDS

<222> (369)..(1158)

<220>

<221> mat_peptide

<222> (94)..()

<400> 3

atg	ctc	ctc	cta	cca	ctc	ctc	tca	gcc	gtc	act	ctc	gcg	gta	gca	agt	48
Met	Leu	Leu	Leu	Pro	Leu	Leu	Ser	Ala	Val	Thr	Leu	Ala	Val	Ala	Ser	
	-30					-25					-20					

cca	cta	gct	tcc	gtc	gaa	gag	tac	gcc	aag	tct	ctc	gaa	gac	aga	g	94
Pro	Leu	Ala	Ser	Val	Glu	Glu	Tyr	Ala	Lys	Ser	Leu	Glu	Asp	Arg		
-15				-10					-5				-1			

gtaagcacca aactctcctc catatcatgc tatatactca tcacactccc ag ct gtg															151	
															Ala Val	
															1	

act	gtg	tct	tcg	tca	gac	tac	aac	aac	ttc	aag	ttc	tac	atc	caa	cat	199
Thr	Val	Ser	Ser	Ser	Asp	Tyr	Asn	Asn	Phe	Lys	Phe	Tyr	Ile	Gln	His	
	5					10					15					

ggc	gcc	gca	gca	tac	tgt	aac	tcc	gaa	gcc	tca	gct	ggc	gca	aag	atc	247
Gly	Ala	Ala	Ala	Tyr	Cys	Asn	Ser	Glu	Ala	Ser	Ala	Gly	Ala	Lys	Ile	
20					25					30						

acc	tgc	gca	agc	aac	ggt	tgt	cca	acc	gtc	cag	tcc	aac	ggc	gca	acc	295
Thr	Cys	Ala	Ser	Asn	Gly	Cys	Pro	Thr	Val	Gln	Ser	Asn	Gly	Ala	Thr	
35				40					45				50			

atc	gtg	gca	tcc	ttc	ct	gtaagtcacg cccagtcaca aacatctcat										342
Ile	Val	Ala	Ser	Phe	Leu											
				55												

acctcactat tatatgactt cttcag t ggt tcc aag act ggc atc ggc ggt															393	
															Gly Ser Lys Thr Gly Ile Gly Gly	
															60	

tat	gtc	gca	aca	gat	tca	tcc	cgc	aag	gaa	atc	gtc	gtc	tcg	atc	cgt	441
Tyr	Val	Ala	Thr	Asp	Ser	Ser	Arg	Lys	Glu	Ile	Val	Val	Ser	Ile	Arg	
65				70					75					80		

gga	agc	agc	aac	atc	cgc	aac	tgg	ctt	aca	aac	ctc	gac	ttt	gac	cag	489
Gly	Ser	Ser	Asn	Ile	Arg	Asn	Trp	Leu	Thr	Asn	Leu	Asp	Phe	Asp	Gln	
			85					90					95			

tcc	gac	tgc	agc	ttg	acc	tcc	ggc	tgc	ggc	gta	cac	tcg	ggc	ttc	cag	537
Ser	Asp	Cys	Ser	Leu	Thr	Ser	Gly	Cys	Gly	Val	His	Ser	Gly	Phe	Gln	
			100				105						110			

aac gcc tgg gac gag atc tcg gag aga gcc act gct gct gtg gcc aag															585	
															Page 6	

10055-WO.ST25

Asn	Ala	Trp	Asp	Glu	Ile	Ser	Glu	Arg	Ala	Thr	Ala	Ala	Val	Ala	Lys	
		115					120					125				
gca	cgc	aag	gca	aac	tct	ggt	ttc	aag	gtc	att	gct	aca	ggc	cac	tcc	633
Ala	Arg	Lys	Ala	Asn	Ser	Gly	Phe	Lys	Val	Ile	Ala	Thr	Gly	His	Ser	
		130				135					140					
ctc	ggt	ggt	gcg	gtc	gct	aca	ttg	gct	gct	gcg	aat	ctg	agg	gtt	ggt	681
Leu	Gly	Gly	Ala	Val	Ala	Thr	Leu	Ala	Ala	Gly	Asn	Leu	Arg	Val	Gly	
					150					155					160	
ggc	aca	ccc	gtg	gac	atc	tac	acg	tac	ggt	gct	cct	cga	gtg	ggc	aac	729
Gly	Thr	Pro	Val	Asp	Ile	Tyr	Thr	Tyr	Gly	Ala	Pro	Arg	Val	Gly	Asn	
				165					170					175		
gcc	cag	ctt	tca	gcg	ttc	atc	tcg	aac	caa	gct	ggc	ggg	gaa	tat	cgc	777
Ala	Gln	Leu	Ser	Ala	Phe	Ile	Ser	Asn	Gln	Ala	Gly	Gly	Glu	Tyr	Arg	
			180					185					190			
gtt	act	cac	gcc	aga	gac	ccc	gtg	cct	cgt	ctg	ccc	cct	ctg	gtg	ttt	825
Val	Thr	His	Ala	Arg	Asp	Pro	Val	Pro	Arg	Leu	Pro	Pro	Leu	Val	Phe	
		195					200					205				
gga	tac	agg	cac	act	tcg	ccc	gag	tac	tgg	cta	tct	ggc	ggc	ggt	ggc	873
Gly	Tyr	Arg	His	Thr	Ser	Pro	Glu	Tyr	Trp	Leu	Ser	Gly	Gly	Gly	Gly	
		210				215					220					
gac	aag	att	gat	tat	acc	atc	agc	gat	atc	aag	gtc	tgt	gag	ggc	gcc	921
Asp	Lys	Ile	Asp	Tyr	Thr	Ile	Ser	Asp	Ile	Lys	Val	Cys	Glu	Gly	Ala	
					230					235					240	
gct	aat	ctc	cag	tgt	aac	ggt	ggc	acg	ctg	ggt	ttg	gac	att	gcg	gct	969
Ala	Asn	Leu	Gln	Cys	Asn	Gly	Gly	Thr	Leu	Gly	Leu	Asp	Ile	Ala	Ala	
				245					250					255		
cat	ctg	cat	tac	ttc	cag	cac	act	gat	gct	tgc	tcg	gcg	gga	ggc	att	1017
His	Leu	His	Tyr	Phe	Gln	His	Thr	Asp	Ala	Cys	Ser	Ala	Gly	Gly	Ile	
			260					265					270			
tct	ttt	aga	cga	tac	agg	agt	gct	aag	cgt	gaa	ggt	atc	gcc	aag	agg	1065
Ser	Phe	Arg	Arg	Tyr	Arg	Ser	Ala	Lys	Arg	Glu	Gly	Ile	Ala	Lys	Arg	
		275					280					285				
gct	gat	atg	tcg	gat	gct	gag	ctg	gag	aag	aag	ctc	aac	tct	tat	gtt	1113
Ala	Asp	Met	Ser	Asp	Ala	Glu	Leu	Glu	Lys	Lys	Leu	Asn	Ser	Tyr	Val	
						295					300					
gag	atg	gat	aag	gag	tat	gtg	gat	agc	cat	aag	aat	cgt	tca	tca	taa	1161
Glu	Met	Asp	Lys	Glu	Tyr	Val	Asp	Ser	His	Lys	Asn	Arg	Ser	Ser		
					310					315						

<210> 4

<211> 350

<212> PRT

<213> Fusarium sulphureum

<400> 4

Met Leu Leu Leu Pro Leu Leu Ser Ala Val Thr Leu Ala Val Ala Ser
 -30 -25 -20

Pro Leu Ala Ser Val Glu Glu Tyr Ala Lys Ser Leu Glu Asp Arg Ala

10055-WO.ST25																			
-15					-10					-5					-1 1				
Val	Thr	Val	Ser	Ser	Ser	Asp	Tyr	Asn	Asn	Phe	Lys	Phe	Tyr	Ile	Gln				
			5					10					15						
His	Gly	Ala	Ala	Ala	Tyr	Cys	Asn	Ser	Glu	Ala	Ser	Ala	Gly	Ala	Lys				
		20					25					30							
Ile	Thr	Cys	Ala	Ser	Asn	Gly	Cys	Pro	Thr	Val	Gln	Ser	Asn	Gly	Ala				
	35					40					45								
Thr	Ile	Val	Ala	Ser	Phe	Leu	Gly	Ser	Lys	Thr	Gly	Ile	Gly	Gly	Tyr				
50					55					60					65				
Val	Ala	Thr	Asp	Ser	Ser	Arg	Lys	Glu	Ile	Val	Val	Ser	Ile	Arg	Gly				
				70					75					80					
Ser	Ser	Asn	Ile	Arg	Asn	Trp	Leu	Thr	Asn	Leu	Asp	Phe	Asp	Gln	Ser				
			85					90					95						
Asp	Cys	Ser	Leu	Thr	Ser	Gly	Cys	Gly	Val	His	Ser	Gly	Phe	Gln	Asn				
		100					105					110							
Ala	Trp	Asp	Glu	Ile	Ser	Glu	Arg	Ala	Thr	Ala	Ala	Val	Ala	Lys	Ala				
		115				120					125								
Arg	Lys	Ala	Asn	Ser	Gly	Phe	Lys	Val	Ile	Ala	Thr	Gly	His	Ser	Leu				
130					135					140					145				
Gly	Gly	Ala	Val	Ala	Thr	Leu	Ala	Ala	Ala	Asn	Leu	Arg	Val	Gly	Gly				
				150					155					160					
Thr	Pro	Val	Asp	Ile	Tyr	Thr	Tyr	Gly	Ala	Pro	Arg	Val	Gly	Asn	Ala				
			165					170					175						
Gln	Leu	Ser	Ala	Phe	Ile	Ser	Asn	Gln	Ala	Gly	Gly	Glu	Tyr	Arg	Val				
		180					185					190							
Thr	His	Ala	Arg	Asp	Pro	Val	Pro	Arg	Leu	Pro	Pro	Leu	Val	Phe	Gly				
	195					200					205								
Tyr	Arg	His	Thr	Ser	Pro	Glu	Tyr	Trp	Leu	Ser	Gly	Gly	Gly	Gly	Asp				
210					215					220					225				
Lys	Ile	Asp	Tyr	Thr	Ile	Ser	Asp	Ile	Lys	Val	Cys	Glu	Gly	Ala	Ala				
				230					235					240					
Asn	Leu	Gln	Cys	Asn	Gly	Gly	Thr	Leu	Gly	Leu	Asp	Ile	Ala	Ala	His				
			245					250					255						
Leu	His	Tyr	Phe	Gln	His	Thr	Asp	Ala	Cys	Ser	Ala	Gly	Gly	Ile	Ser				
		260					265					270							

10055-WO.ST25

Phe Arg Arg Tyr Arg Ser Ala Lys Arg Glu Gly Ile Ala Lys Arg Ala
 275 280 285

Asp Met Ser Asp Ala Glu Leu Glu Lys Lys Leu Asn Ser Tyr Val Glu
 290 295 300 305

Met Asp Lys Glu Tyr Val Asp Ser His Lys Asn Arg Ser Ser
 310 315

<210> 5

<211> 1103

<212> DNA

<213> Acremonium berkeleyanum

<220>

<221> CDS

<222> (1)..(85)

<220>

<221> CDS

<222> (141)..(306)

<220>

<221> CDS

<222> (365)..(1100)

<220>

<221> mat_peptide

<222> (85)..()

<400> 5

atg ctc gcc cta tcc ctt ctt tct att gct gcc ctg gcg gta gct agt 48
 Met Leu Ala Leu Ser Leu Leu Ser Ile Ala Ala Leu Ala Val Ala Ser
 -25 -20 -15

ccc ttg gcc gat tat tcc aag gcc ttg gaa gac cga g gtgagtccaa 95
 Pro Leu Ala Asp Tyr Ser Lys Ala Leu Glu Asp Arg
 -10 -5 -1

ccacttacac acctccaggc agaaacgatt gctaagtaaa cacag cc atc tcg gtc 151
 Ala Ile Ser Val
 1

10055-WO.ST25															
acc gat gga gac ctg aac aac ttc aag ttc tac gtc caa cac gct gcc	Thr Asp Gly Asp Leu Asn Asn Phe Lys Phe Tyr Val Gln His Ala Ala	5	10	15	20	25	30	35	40	45	50	55	60	65	
199															
gcc gca tac tgc aat gtc aac act cca gcg ggt caa gct gtg aag tgc	Ala Ala Tyr Cys Asn Val Asn Thr Pro Ala Gly Gln Ala Val Lys Cys	25	30	35	40	45	50	55	60	65	70	75	80	85	247
ggc gga tcc aca tgc tct gct gtc gaa ggc gac agg gtg aca gtc gtc	Gly Gly Ser Thr Cys Ser Ala Val Glu Gly Asp Arg Val Thr Val Val	40	45	50	55	60	65	70	75	80	85	90	95	100	295
gca tcc ttc aa gtacgttcct cttctatgtt agacttaaag aagcaatccc	Ala Ser Phe Asn	55	60	65	70	75	80	85	90	95	100	105	110	115	346
agctaacgtc cccatcag c ggc gct gga aca gga att ggc ggc tat gta gcc	Gly Ala Gly Thr Gly Ile Gly Gly Tyr Val Ala	60	65	70	75	80	85	90	95	100	105	110	115	120	398
acc gac aac gcc cgc tct gag atc gtt gtc tcc atc cgc ggt agt agc	Thr Asp Asn Ala Arg Ser Glu Ile Val Val Ser Ile Arg Gly Ser Ser	70	75	80	85	90	95	100	105	110	115	120	125	130	446
aat atc cgc aac tgg att gcc aac ata gag ttt gcg cag cag gac tgc	Asn Ile Arg Asn Trp Ile Ala Asn Ile Glu Phe Ala Gln Gln Asp Cys	85	90	95	100	105	110	115	120	125	130	135	140	145	494
tcc ctt gtt gct ggc tgc ggt gtg cac act ggc ttc cag aag gca tgg	Ser Leu Val Ala Gly Cys Gly Val His Thr Gly Phe Gln Lys Ala Trp	100	105	110	115	120	125	130	135	140	145	150	155	160	542
aac gag atc tcc gcc aac gtc aag gcc gct gtg gca tct gcg aag cag	Asn Glu Ile Ser Ala Asn Val Lys Ala Ala Val Ala Ser Ala Lys Gln	120	125	130	135	140	145	150	155	160	165	170	175	180	590
gcg aac cca agc tac aag atc atc tcg act ggt cac tcc ctc ggc ggt	Ala Asn Pro Ser Tyr Lys Ile Ile Ser Thr Gly His Ser Leu Gly Gly	135	140	145	150	155	160	165	170	175	180	185	190	195	638
gcg gtg gct acc ctc gcg gcc gcg tac ctg cgt aag gat ggc aat gct	Ala Val Ala Thr Leu Ala Ala Ala Tyr Leu Arg Lys Asp Gly Asn Ala	150	155	160	165	170	175	180	185	190	195	200	205	210	686
gtc gat ctg tac aca tac ggc tcg cca cga gtc ggc aac ggt gtc ttc	Val Asp Leu Tyr Thr Tyr Gly Ser Pro Arg Val Gly Asn Gly Val Phe	165	170	175	180	185	190	195	200	205	210	215	220	225	734
tct aac ttc gtc agc caa caa gcc gcc tct gaa ttc cgc gtc acc cac	Ser Asn Phe Val Ser Gln Gln Ala Gly Ser Glu Phe Arg Val Thr His	180	185	190	195	200	205	210	215	220	225	230	235	240	782
ggc gac gac ccc gtc ccc cgt ctg ccc cca atc gtc ttc ggc tac cgc	Gly Asp Asp Pro Val Pro Arg Leu Pro Pro Ile Val Phe Gly Tyr Arg	200	205	210	215	220	225	230	235	240	245	250	255	260	830
cac acc acc ccg gag tac tgg ctc gac ggc ggc tcc ctg gac gtg acg	His Thr Thr Pro Glu Tyr Trp Leu Asp Gly Gly Ser Leu Asp Val Thr	215	220	225	230	235	240	245	250	255	260	265	270	275	878
tac aac ctc gac gag atc aag gtg tgc gag ggc aat gcg aat gtg aac	Tyr Asn Leu Asp Glu Ile Lys Val Cys Glu Gly Asn Ala Asn Val Asn	230	235	240	245	250	255	260	265	270	275	280	285	290	926
tgc aac ggt ggt acg ttt ggc ctt gat atc ctt gcg cat ttg cgg tac	Cys Asn Gly Gly Thr Phe Gly Leu Asp Ile Leu Ala His Leu Arg Tyr	245	250	255	260	265	270	275	280	285	290	295	300	305	974
cta cag gat gta tca ggt tgt gcg ccg atc ggc atc ttc tgg aag cgc		305	310	315	320	325	330	335	340	345	350	355	360	365	1022

10055-WO.ST25

Leu Gln Asp Val Ser Gly Cys Ala Pro Ile Gly Ile Phe Trp Lys Arg
 260 265 270 275

gag gag atg tcg gat gaa gag ttg gag aag aag gtg aat gag tat gtc 1070
 Glu Glu Met Ser Asp Glu Glu Leu Glu Lys Lys Val Asn Glu Tyr Val
 280 285 290

cag gcc gac aag gac ttt gtg gct ggg ttg tag 1103
 Gln Ala Asp Lys Asp Phe Val Ala Gly Leu
 295 300

<210> 6

<211> 329

<212> PRT

<213> Acremonium berkeleyanum

<400> 6

Met Leu Ala Leu Ser Leu Leu Ser Ile Ala Ala Leu Ala Val Ala Ser
 -25 -20 -15

Pro Leu Ala Asp Tyr Ser Lys Ala Leu Glu Asp Arg Ala Ile Ser Val
 -10 -5 -1 1

Thr Asp Gly Asp Leu Asn Asn Phe Lys Phe Tyr Val Gln His Ala Ala
 5 10 15 20

Ala Ala Tyr Cys Asn Val Asn Thr Pro Ala Gly Gln Ala Val Lys Cys
 25 30 35

Gly Gly Ser Thr Cys Ser Ala Val Glu Gly Asp Arg Val Thr Val Val
 40 45 50

Ala Ser Phe Asn Gly Ala Gly Thr Gly Ile Gly Gly Tyr Val Ala Thr
 55 60 65

Asp Asn Ala Arg Ser Glu Ile Val Val Ser Ile Arg Gly Ser Ser Asn
 70 75 80

Ile Arg Asn Trp Ile Ala Asn Ile Glu Phe Ala Gln Gln Asp Cys Ser
 85 90 95 100

Leu Val Ala Gly Cys Gly Val His Thr Gly Phe Gln Lys Ala Trp Asn
 105 110 115

Glu Ile Ser Ala Asn Val Lys Ala Ala Val Ala Ser Ala Lys Gln Ala
 120 125 130

Asn Pro Ser Tyr Lys Ile Ile Ser Thr Gly His Ser Leu Gly Gly Ala
 135 140 145

Val Ala Thr Leu Ala Ala Ala Tyr Leu Arg Lys Asp Gly Asn Ala Val
 Page 11

150											155	10055-WO.ST25										160			
Asp 165	Leu	Tyr	Thr	Tyr	Gly 170	Ser	Pro	Arg	Val	Gly 175	Asn	Gly	Val	Phe	Ser 180										
Asn	Phe	Val	Ser	Gln 185	Gln	Ala	Gly	Ser	Glu 190	Phe	Arg	Val	Thr	His 195	Gly										
Asp	Asp	Pro	Val 200	Pro	Arg	Leu	Pro	Pro 205	Ile	Val	Phe	Gly	Tyr 210	Arg	His										
Thr	Thr	Pro 215	Glu	Tyr	Trp	Leu	Asp 220	Gly	Gly	Ser	Leu	Asp 225	Val	Thr	Tyr										
Asn	Leu 230	Asp	Glu	Ile	Lys	Val 235	Cys	Glu	Gly	Asn	Ala 240	Asn	Val	Asn	Cys										
Asn 245	Gly	Gly	Thr	Phe	Gly 250	Leu	Asp	Ile	Leu	Ala 255	His	Leu	Arg	Tyr	Leu 260										
Gln	Asp	Val	Ser	Gly 265	Cys	Ala	Pro	Ile	Gly 270	Ile	Phe	Trp	Lys	Arg 275	Glu										
Glu	Met	Ser	Asp 280	Glu	Glu	Leu	Glu	Lys 285	Lys	Val	Asn	Glu	Tyr 290	Val	Gln										
Ala	Asp	Lys 295	Asp	Phe	Val	Ala	Gly 300	Leu																	

<210>	7
<211>	1159
<212>	DNA
<213>	<i>Fusarium culmorum</i>

```
<220>
<221> CDS
<222> (1)..(91)
```

```
<220>
<221> CDS
<222> (144)..(309)
```

<220>
<221> CDS

10055-WO.ST25

<222> (367)..(1156)

<220>

<221> mat_peptide

<222> (91)..()

<400> 7

atg cgt ctc ctg tca ctc ctc tca gtt gtc acc ctt gtg gta gcc agc	48
Met Arg Leu Leu Ser Leu Leu Ser Val Val Thr Leu Val Val Ala Ser	
-30 -25 -20 -15	
cct ctg agc gtt gaa gaa tac gcc aag gct ctc gat gaa cga g	91
Pro Leu Ser Val Glu Glu Tyr Ala Lys Ala Leu Asp Glu Arg	
-10 -5 -1	
gtacacacga tcatcattcc tataacaaac tccataactca caaagtctct ag ct gtc	148
	Ala Val
	1
tct gtc tcc acc acc gac ttt ggc aat ttc aag ttc tac atc cag cac	196
Ser Val Ser Thr Thr Asp Phe Gly Asn Phe Lys Phe Tyr Ile Gln His	
5 10 15	
ggc gcc gca gca tac tgc aac tcc gaa gcc ccg gcc ggt gca aag gtc	244
Gly Ala Ala Ala Tyr Cys Asn Ser Glu Ala Pro Ala Gly Ala Lys Val	
20 25 30	
acc tgc agc gga aac ggc tgt cca act gtt cag tcc aac ggt gtt acc	292
Thr Cys Ser Gly Asn Gly Cys Pro Thr Val Gln Ser Asn Gly Val Thr	
35 40 45 50	
atc gtg gca tcc ttc ac gtaagtccca cccggcacia acacatcacc	339
Ile Val Ala Ser Phe Thr	
55	
aactccaagc ttacacgttt ctctcag t gga tcc aag act gga atc ggc ggc	391
	Gly Ser Lys Thr Gly Ile Gly Gly
	60
tac gtc gct aca gac cct aca cgc aag gag atc gtc gtc tcg ttc cgt	439
Tyr Val Ala Thr Asp Pro Thr Arg Lys Glu Ile Val Val Ser Phe Arg	
65 70 75 80	
ggt agc atc aac atc cgc aac tgg ctt acc aac ctc gac ttc gac cag	487
Gly Ser Ile Asn Ile Arg Asn Trp Leu Thr Asn Leu Asp Phe Asp Gln	
85 90 95	
gac gac tgc agc ctg acc tcg ggc tgt ggt gtt cac tca ggc ttc cag	535
Asp Asp Cys Ser Leu Thr Ser Gly Cys Gly Val His Ser Gly Phe Gln	
100 105 110	
aaa gcc tgg aac gag atc tca gcc gcg gca acc gcc gct gtc gca aag	583
Lys Ala Trp Asn Glu Ile Ser Ala Ala Ala Thr Ala Ala Val Ala Lys	
115 120 125	
gcc cgc aag gca aac cct tcg ttc aag gtc gtc tcc gta ggt cac tcc	631
Ala Arg Lys Ala Asn Pro Ser Phe Lys Val Val Ser Val Gly His Ser	
130 135 140	
ctg ggt ggt gct gta gct aca ctg gca ggc gcg aac cta cga gtt ggt	679
Leu Gly Gly Ala Val Ala Thr Leu Ala Gly Ala Asn Leu Arg Val Gly	
145 150 155 160	

10055-WO.ST25

gga aca ccc ctt gac att tac acc tac ggt tca ccc cga gtt gga aac	727
Gly Thr Pro Leu Asp 165 Ile Tyr Thr Tyr Gly Ser Pro Arg Val Gly Asn 175	
aca cag ctc gct gct ttt gtc tcg aac cag gct ggt gga gag ttc cgc	775
Thr Gln Leu Ala 180 Phe Val Ser Asn 185 Gln Ala Gly Gly Glu Phe Arg 190	
gtt acg aac gcc aaa gac ccc gtg cct cgt ctc ccc cct ctg atc ttt	823
Val Thr Asn 195 Ala Lys Asp Pro 200 Val Pro Arg Leu Pro 205 Leu Ile Phe	
gga tac cga cac aca tcc ccc gag tac tgg ctg tct ggc agc gga ggt	871
Gly Tyr Arg His Thr Ser Pro 215 Glu Tyr Trp Leu Ser 220 Gly Ser Gly Gly	
gac aag atc gac tac acc atc aac gat gtc aag gtc tgt gaa ggt gcc	919
Asp Lys Ile Asp Tyr 230 Thr Ile Asn Asp Val 235 Lys Val Cys Glu Gly Ala 240	
gcc aac ctc cag tgc aac ggt gga aca ctc gga ttg gat atc gat gcc	967
Ala Asn Leu Gln Cys 245 Asn Gly Gly Thr Leu Gly Leu Asp Ile Asp Ala 255	
cat ctc cac tac ttc cag gca act gat gct tgc tct gct ggc ggc atc	1015
His Leu His Tyr 260 Phe Gln Ala Thr Asp 265 Ala Cys Ser Ala Gly Gly Ile 270	
tcg tgg aga aga tac agg agc gcc aag cgt gag agc atc tca gag agg	1063
Ser Trp Arg 275 Arg Tyr Arg Ser Ala 280 Lys Arg Glu Ser Ile 285 Ser Glu Arg	
gct acc atg acc gac gcc gag ctc gag aag aag ctc aac agc tat gtt	1111
Ala Thr Met Thr Asp Ala 295 Glu Leu Glu Lys Lys 300 Leu Asn Ser Tyr Val	
gag atg gat aag gag tat atc aag act cac gcc agc cgc tca tca tag	1159
Glu Met Asp Lys Glu Tyr 310 Ile Lys Thr His Ala 315 Ser Arg Ser Ser	

<210> 8

<211> 349

<212> PRT

<213> Fusarium culmorum

<400> 8

Met Arg Leu Leu Ser Leu Leu Ser Val Val Thr Leu Val Val Ala Ser
-30 -25 -20 -15

Pro Leu Ser Val Glu Glu Tyr Ala Lys Ala Leu Asp Glu Arg Ala Val
-10 -5 -1 1

Ser Val Ser Thr Thr Asp Phe Gly Asn Phe Lys Phe Tyr Ile Gln His
5 10 15

Gly Ala Ala Ala Tyr Cys Asn Ser Glu Ala Pro Ala Gly Ala Lys Val
20 25 30

10055-WO.ST25

Thr Cys Ser Gly Asn Gly Cys Pro Thr Val Gln Ser Asn Gly Val Thr
 35 40 45 50

Ile Val Ala Ser Phe Thr Gly Ser Lys Thr Gly Ile Gly Gly Tyr Val
 55 60 65

Ala Thr Asp Pro Thr Arg Lys Glu Ile Val Val Ser Phe Arg Gly Ser
 70 75 80

Ile Asn Ile Arg Asn Trp Leu Thr Asn Leu Asp Phe Asp Gln Asp Asp
 85 90 95

Cys Ser Leu Thr Ser Gly Cys Gly Val His Ser Gly Phe Gln Lys Ala
 100 105 110

Trp Asn Glu Ile Ser Ala Ala Ala Thr Ala Ala Val Ala Lys Ala Arg
 115 120 125 130

Lys Ala Asn Pro Ser Phe Lys Val Val Ser Val Gly His Ser Leu Gly
 135 140 145

Gly Ala Val Ala Thr Leu Ala Gly Ala Asn Leu Arg Val Gly Gly Thr
 150 155 160

Pro Leu Asp Ile Tyr Thr Tyr Gly Ser Pro Arg Val Gly Asn Thr Gln
 165 170 175

Leu Ala Ala Phe Val Ser Asn Gln Ala Gly Gly Glu Phe Arg Val Thr
 180 185 190

Asn Ala Lys Asp Pro Val Pro Arg Leu Pro Pro Leu Ile Phe Gly Tyr
 195 200 205 210

Arg His Thr Ser Pro Glu Tyr Trp Leu Ser Gly Ser Gly Gly Asp Lys
 215 220 225

Ile Asp Tyr Thr Ile Asn Asp Val Lys Val Cys Glu Gly Ala Ala Asn
 230 235 240

Leu Gln Cys Asn Gly Gly Thr Leu Gly Leu Asp Ile Asp Ala His Leu
 245 250 255

His Tyr Phe Gln Ala Thr Asp Ala Cys Ser Ala Gly Gly Ile Ser Trp
 260 265 270

Arg Arg Tyr Arg Ser Ala Lys Arg Glu Ser Ile Ser Glu Arg Ala Thr
 275 280 285 290

Met Thr Asp Ala Glu Leu Glu Lys Lys Leu Asn Ser Tyr Val Glu Met
 295 300 305

310

10055-WO.ST25
315

<210> 9
 <211> 1150
 <212> DNA
 <213> Fusarium solani

<220>
 <221> CDS
 <222> (1)..(93)

<220>
 <221> CDS
 <222> (149)..(316)

<220>
 <221> CDS
 <222> (365)..(1147)

<220>
 <221> mat_peptide
 <222> (149)..()

<400> 9
 atg cgc ctg ctt cct ctc ctc tcg gtc gta acg ctc act gcg gcg agt 48
 Met Arg Leu Leu Pro Leu Leu Ser Val Val Thr Leu Thr Ala Ala Ser
 -30 -25 -20
 cct ata gcc tcc gtc cag gag tac act gac gcc ttg gag aag aga 93
 Pro Ile Ala Ser Val Gln Glu Tyr Thr Asp Ala Leu Glu Lys Arg
 -15 -10 -5 -1
 ggtaaccacc aacactcccc taagacagac ctgcacctaa caagtaaact ctagt gct 151
 Ala
 1
 atc acc gcc tct caa ctt gac tat gaa aac ttc aag ttt tac atc cag 199
 Ile Thr Ala Ser Gln Leu Asp Tyr Glu Asn Phe Lys Phe Tyr Ile Gln
 5 10 15
 cac ggt gcc gca gcg tat tgc aac tct gag acg gcc tct ggt caa aaa 247
 His Gly Ala Ala Ala Tyr Cys Asn Ser Glu Thr Ala Ser Gly Gln Lys
 20 25 30
 ata acc tgc aac gac aac ggc tgc aaa ggc atc gag gcc aac aac gcc 295
 Ile Thr Cys Asn Asp Asn Gly Cys Lys Gly Ile Glu Ala Asn Asn Ala
 35 40 45

10055-WO.ST25

ata atc gta gca tcc ttc gtg taagctcccc tttccctca cggaaccctt Ile Ile Val Ala Ser Phe Val 50 55	346
caactgacac gtagcagc ggc acg ggc act ggc atc gga ggc tac gtc tcc Gly Thr Gly Thr Gly Ile Gly Gly Tyr Val Ser 60 65	397
act gac aat gtc cgt aag gag att gtc ctc tcg att cgc ggc agc agc Thr Asp Asn Val Arg Lys Glu Ile Val Leu Ser Ile Arg Gly Ser Ser 70 75 80	445
aac atc cgc aac tgg ctc acc aac gtc gac ttt ggc cag tcc agc tgc Asn Ile Arg Asn Trp Leu Thr Asn Val Asp Phe Gly Gln Ser Ser Cys 85 90 95	493
tcc tac gtc cgc gac tgc gga gtc cac acg ggc ttc cgc aat gcc tgg Ser Tyr Val Arg Asp Cys Gly Val His Thr Gly Phe Arg Asn Ala Trp 100 105 110 115	541
gat gag att gcc cag cgc gcg agg gac gcc gtt gcc aag gcc cgc gcc Asp Glu Ile Ala Gln Arg Ala Arg Asp Ala Val Ala Lys Ala Arg Ala 120 125 130	589
atg aac ccg tcc tac aag gtc atc tcc acg ggc cac tct ctc ggc ggt Met Asn Pro Ser Tyr Lys Val Ile tcc Thr Gly His Ser Leu Gly Gly 135 140 145	637
gct gtc gca act ctg ggt gcc gct gac ctg agg tcc aag gga acc gca Ala Val Ala Thr Leu Gly Ala Ala 155 Asp Leu Arg Ser Lys Gly Thr Ala 150 155 160	685
gtt gac atc ttc acc ttt ggt gct ccc cgt gta ggc aac gct gaa ctc Val Asp Ile Phe Thr Phe Gly Ala Pro Arg Val Gly Asn Ala Glu Leu 165 170 175	733
tca gca ttc atc acg gcc cag gcc ggc ggc gag ttc cgt gtc act cat Ser Ala Phe Ile Thr Ala Gln Ala Gly Gly Glu Phe Arg Val Thr His 180 185 190 195	781
ggc cgt gat ccc gtg ccc cgt ctg cct ccc atc gtc ttt ggc tac aga Gly Arg Asp Pro Val Pro Arg Leu Pro Pro Ile Val Phe Gly Tyr Arg 200 205 210	829
cac aca tcg ccc gag tac tgg ctg gcc ggc ggt gca tcc acc aag atc His Thr Ser Pro Glu Tyr Trp Leu Ala 220 Gly Gly Ala Ser Thr Lys Ile 215 220 225	877
gac tac tcc gtc aac gac atc aag gtc tgt gaa ggc gcc gcc aat ctc Asp Tyr Ser Val Asn Asp Ile Lys Val Cys Glu Gly Ala Ala Asn Leu 230 235 240	925
gcc tgc aac ggc ggt aca cta ggc ctg gat atc atc gct cat ctg cgc Ala Cys Asn Gly Gly Thr Leu 250 Gly Leu Asp Ile Ile Ala His Leu Arg 245 250 255	973
tac ttc cag aac acc gat gcc tgc aca gcg ggc ggt atc tcg tgg aag Tyr Phe Gln Asn Thr Asp Ala Cys Thr Ala Gly Gly Ile Ser Trp Lys 260 265 270 275	1021
aga ggg gac aag gcc aag cgt gac gag atc ccc aag cgc cag gag ggc Arg Gly Asp Lys Ala Lys Arg Asp Glu Ile Pro Lys Arg Gln Glu Gly 280 285 290	1069
atg acg gat gag gag ttg gag cag aag ctc aac gac tat gtc gcc atg Met Thr Asp Glu Glu Leu Glu Gln Lys Leu Asn Asp Tyr Val Ala Met 295 300 305	1117

10055-WO.ST25
 gac aag gag tac gtg gac agc cat aag atc tag
 Asp Lys Glu Tyr Val Asp Ser His Lys Ile
 310 315

1150

<210> 10

<211> 348

<212> PRT

<213> Fusarium solani

<400> 10

Met Arg Leu Leu Pro Leu Leu Ser Val Val Thr Leu Thr Ala Ala Ser
 -30 -25 -20

Pro Ile Ala Ser Val Gln Glu Tyr Thr Asp Ala Leu Glu Lys Arg Ala
 -15 -10 -5 -1 1

Ile Thr Ala Ser Gln Leu Asp Tyr Glu Asn Phe Lys Phe Tyr Ile Gln
 5 10 15

His Gly Ala Ala Ala Tyr Cys Asn Ser Glu Thr Ala Ser Gly Gln Lys
 20 25 30

Ile Thr Cys Asn Asp Asn Gly Cys Lys Gly Ile Glu Ala Asn Asn Ala
 35 40 45

Ile Ile Val Ala Ser Phe Val Gly Thr Gly Thr Gly Ile Gly Gly Tyr
 50 55 60 65

Val Ser Thr Asp Asn Val Arg Lys Glu Ile Val Leu Ser Ile Arg Gly
 70 75 80

Ser Ser Asn Ile Arg Asn Trp Leu Thr Asn Val Asp Phe Gly Gln Ser
 85 90 95

Ser Cys Ser Tyr Val Arg Asp Cys Gly Val His Thr Gly Phe Arg Asn
 100 105 110

Ala Trp Asp Glu Ile Ala Gln Arg Ala Arg Asp Ala Val Ala Lys Ala
 115 120 125

Arg Ala Met Asn Pro Ser Tyr Lys Val Ile Ser Thr Gly His Ser Leu
 130 135 140 145

Gly Gly Ala Val Ala Thr Leu Gly Ala Ala Asp Leu Arg Ser Lys Gly
 150 155 160

Thr Ala Val Asp Ile Phe Thr Phe Gly Ala Pro Arg Val Gly Asn Ala
 165 170 175

10055-WO.ST25

Glu Leu Ser Ala Phe Ile Thr Ala Gln Ala Gly Gly Glu Phe Arg Val
 180 185 190

Thr His Gly Arg Asp Pro Val Pro Arg Leu Pro Pro Ile Val Phe Gly
 195 200 205

Tyr Arg His Thr Ser Pro Glu Tyr Trp Leu Ala Gly Gly Ala Ser Thr
 210 215 220 225

Lys Ile Asp Tyr Ser Val Asn Asp Ile Lys Val Cys Glu Gly Ala Ala
 230 235 240

Asn Leu Ala Cys Asn Gly Gly Thr Leu Gly Leu Asp Ile Ile Ala His
 245 250 255

Leu Arg Tyr Phe Gln Asn Thr Asp Ala Cys Thr Ala Gly Gly Ile Ser
 260 265 270

Trp Lys Arg Gly Asp Lys Ala Lys Arg Asp Glu Ile Pro Lys Arg Gln
 275 280 285

Glu Gly Met Thr Asp Glu Glu Leu Glu Gln Lys Leu Asn Asp Tyr Val
 290 295 300 305

Ala Met Asp Lys Glu Tyr Val Asp Ser His Lys Ile
 310 315

<210> 11

<211> 35

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> Primer

<400> 11

cagtgaattg gcctcgatgg ccgcggccgc gaatt

35

<210> 12

<211> 35

<212> DNA

<213> Artificial/Unknown

10055-WO.ST25

<220>

<221> misc_feature

<222> ()..()

<223> Primer

<400> 12

aattcgcggc cgcggccatc gaggccaatt cactg

35

<210> 13

<211> 34

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> Primer

<400> 13

cacgaaggaa agacgatggc tttcacggtg tctg

34

<210> 14

<211> 34

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> Primer

<400> 14

cagacaccgt gaaagccatc gtctttcctt cgtg

34

10055-WO.ST25

<210> 15
<211> 46
<212> DNA
<213> Artificial/Unknown

<220>
<221> misc_feature
<222> ()..()
<223> Primer

<400> 15
ctatctcttc accatggtac cttaattaaa taccttggtg gaagcg

46

<210> 16
<211> 46
<212> DNA
<213> Artificial/Unknown

<220>
<221> misc_feature
<222> ()..()
<223> Primer

<400> 16
cgcttccaac aaggtattta attaaggtac catggtgaag agatag

46

<210> 17
<211> 37
<212> DNA
<213> Artificial/Unknown

<220>
<221> misc_feature
<222> ()..()

10055-WO.ST25

<223> Primer

<400> 17
cgttctttgt ctgtcagcat gcatctccta tcaactcc

37

<210> 18

<211> 45

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> O..O

<223> Primer

<400> 18
ccagagtttt tgttatgggt aattaatatc gttactgcgt aaatg

45

<210> 19

<211> 20

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> O..O

<223> 1ip2

<400> 19
aarttytaya thcarcaygg

20

<210> 20

<211> 23

<212> DNA

10055-WO.ST25

<213> Artificial/Unknown

<220>

<221> modified_base

<222> (1)..(23)

<223> i

<220>

<221> misc_feature

<222> ()..()

<223> Primer lip3

<400> 20
carcayggng cngcngcnta ytg

23

<210> 21

<211> 23

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> lip10

<220>

<221> modified_base

<222> (1)..(23)

<223> i

<400> 21
ggntgyggng tncaynnngg ntt

23

10055-WO.ST25

<210> 22
<211> 23
<212> DNA
<213> Artificial/Unknown

<220>
<221> modified_base
<222> (1)..(23)
<223> i

<220>
<221> misc_feature
<222> ()..()
<223> lip11

<400> 22
aanccnnnrt gnacnccrca ncc

23

<210> 23
<211> 20
<212> DNA
<213> Artificial/Unknown

<220>
<221> misc_feature
<222> ()..()
<223> Primer lip15

<220>
<221> modified_base
<222> (1)..(20)
<223> i

10055-WO.ST25

<400> 23
ccnccnarns wrtgncngt

20

<210> 24

<211> 20

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> lip17

<220>

<221> modified_base

<222> (1)..(20)

<223> i

<400> 24
ggrrctntyns crtkngtnac

20

<210> 25

<211> 23

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> lip21

<220>

10055-WO.ST25

<221> modified_base

<222> (1)..(23)

<223> i

<400> 25

tcnswngtyt gnckrtancc raa

23

<210> 26

<211> 44

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> Adaptor L

<400> 26

ctaatacgac tcactatagg gctcgagcgg ccgcccgggc aggt

44

<210> 27

<211> 21

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 27N1long

<400> 27

tggacaaccg ttccttgccg a

21

<210> 28

10055-WO.ST25

<211> 25

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 27c1long

<400> 28

tacacgtacg gtgctcctcg agtgg

25

<210> 29

<211> 20

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 27c2

<400> 29

tatctggcgg cggtggcgac

20

<210> 30

<211> 36

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 27N(Bam)

10055-WO.ST25

<400> 30
cgcggtacca tgctcctcct accactcctc tcagcc 36

<210> 31

<211> 48

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 27C(Sal)

<400> 31
acgcgtcgac ttatgatgaa cgattcttat ggctatccac atactcct 48

<210> 32

<211> 30

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> AcN3

<400> 32
tggatccgcc gcacttcaca gcttgacccg 30

<210> 33

<211> 32

<212> DNA

<213> Artificial/Unknown

10055-WO.ST25

<220>

<221> misc_feature

<222> ()..()

<223> Acc3

<400> 33
cggcaacggt gtctttctcta acttcgtcag cc

32

<210> 34

<211> 38

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 152-N (Bcl)

<400> 34
tattatcatg atcaatgctc gccctatccc ttctttct

38

<210> 35

<211> 36

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 152-C(Xho)

<400> 35
ccgctcgagc tacaaccag ccacaaagtc cttgtc

36

<210> 36

10055-WO.ST25

<211> 26

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 140288

<400> 36
ttgaattcat gggtaataac tgatat

26

<210> 37

<211> 44

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 141222

<400> 37
gggtattgtcc tgcagacggc aatttaacgg cttctgcgaa tcgc

44

<210> 38

<211> 45

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 141223

10055-WO.ST25

<400> 38
ggatgctgtt gactccggaa atttaacggt ttggtcttgc atccc

45

<210> 39

<211> 32

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 142778

<400> 39
aaatcaatct attttcaatt caattcatca tt

32

<210> 40

<211> 31

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 142779

<400> 40
ttgaattgaa aatagattga tttaaaactt c

31

<210> 41

<211> 25

<212> DNA

<213> Artificial/Unknown

10055-WO.ST25

<220>

<221> misc_feature

<222> ()..()

<223> 142780

<400> 41

ttgcatgcgt aatcatgggc atagc

25

<210> 42

<211> 30

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 12N1

<400> 42

actgaacagt tggacagccg tttccgctgc

30

<210> 43

<211> 32

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 12C2

<400> 43

ccaggctggg ggagagttcc gcgttacgaa cg

32

10055-WO.ST25

<210> 44

<211> 33

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 12-N (Bcl)

<400> 44

ttgtctgtga tcatgcgtct cctgtcactc etc

33

<210> 45

<211> 44

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 12-C (Sal)

<400> 45

ttagtgcgta aacagctgac tatgatgagc ggctggcgtg agtc

44

<210> 46

<211> 23

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 161000j1

10055-WO.ST25

<400> 46
acaggccact cccttgagg ngc

23

<210> 47

<211> 24

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 161000j2

<400> 47
aggagggaga cgggggacng grtc

24

<210> 48

<211> 28

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 071200j1

<400> 48
gctgaactct cagcattcat cacggccc

28

<210> 49

<211> 28

<212> DNA

<213> Artificial/Unknown

10055-WO.ST25

<220>

<221> misc_feature

<222> ()..()

<223> 071200j2

<400> 49
ccaaaggtga agatgtcaac tgcggttc

28

<210> 50

<211> 26

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 221200j1

<400> 50
cggcggcgag ttccgtgtca ctcacg

26

<210> 51

<211> 25

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 221200j2

<400> 51
gacctcaggt cagcggcacc cagag

25

10055-WO.ST25

<210> 52
<211> 28
<212> DNA
<213> Artificial/Unknown

<220>
<221> misc_feature
<222> ()..()
<223> 170101111

<400> 52
cttgaagttt tcatagtcaa gttgagag

28

<210> 53
<211> 39
<212> DNA
<213> Artificial/Unknown

<220>
<221> misc_feature
<222> ()..()
<223> 290101j2

<400> 53
actagcctcg agctagatct tatggctgtc cacgtactc

39

<210> 54
<211> 45
<212> DNA
<213> Artificial/Unknown

<220>
<221> misc_feature
<222> ()..()

10055-WO.ST25

<223> 020301j1

<400> 54
gcgcgcggat ccaccatgcg cctgcttcct ctctctcgg tcgta

45

<210> 55

<211> 21

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 19670

<400> 55
ccccatcctt taactatagc g

21

<210> 56

<211> 24

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 19671

<400> 56
ctcccttctc tgaacaataa accc

24

<210> 57

<211> 11

<212> DNA

<213> Artificial/Unknown

10055-WO.ST25

<220>

<221> misc_feature

<222> ()..()

<223> nucleotides 134-144

<400> 57
gtactaaaac c

11

<210> 58

<211> 11

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 134-144 altered

<400> 58
ccgttaaatt t

11

<210> 59

<211> 0

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> nucleotides 423-436

<400> 59
000

10055-WO.ST25

<210> 60

<211> 14

<212> DNA

<213> Artificial/Unknown

<220>

<221> misc_feature

<222> ()..()

<223> 423-436 altered

<400> 60
cggcaattta acgg

14

